

BG

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 December 2001 (13.12.2001)

PCT

(10) International Publication Number
WO 01/94587 A2(51) International Patent Classification⁷: C12N 15/16,
15/19, C07K 14/52, 14/575, 16/24, 16/26, C12Q 1/68,
A61K 38/19, 38/22, G01N 33/53

(21) International Application Number: PCT/US01/18476

(22) International Filing Date: 6 June 2001 (06.06.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/210,233 6 June 2000 (06.06.2000) US
60/213,465 23 June 2000 (23.06.2000) US
60/249,019 14 November 2000 (14.11.2000) US(71) Applicant (for all designated States except US): INCYTE
GENOMICS, INC. [US/US]; 3160 Porter Drive, Palo
Alto, CA 94304 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LAL, Preeti
[IN/US]; P.O. Box 5142, Santa Clara, CA 95056 (US).
YUE, Henry [US/US]; 826 Lois Avenue, Sunnyvale,
CA 94087 (US). HE, Ann [CN/US]; 4601 Catalina
Drive, San Jose, CA 95129 (US). NGUYEN, Danniell, B.
[US/US]; 1403 Ridgewood Drive, San Jose, CA 95118
(US). WALIA, Narinder [US/US]; 890 Davis Street #205,
San Leandro, CA 94577 (US). GANDHI, Ameena; R.
[US/US]; 837 Roble Avenue, #1, Menlo Park, CA 94025
(US). AZIMZAI, Yalda [US/US]; 5518 Boulder Canyon
Drive, Castro Valley, CA 94552 (US). BANDMAN, Olga
[US/US]; 366 Anna Avenue, Mountain View, CA 94043
(US). TANG, Y., Tom [US/US]; 4230 Ranwick Court, SanJose, CA 95118 (US). LU, Yan [US/US]; 3885 Corrina
Way, Palo Alto, CA 94303 (US). BAUGHN, Mariah, R.
[US/US]; 14244 Santiago Road, San Leandro, CA 94577
(US). DUGGAN, Brendan, M. [AU/US]; 243 Buena
Vista, #306, Sunnyvale, CA 94086 (US). LEE, Sally
[US/US]; 825 East Evelyn, #425, Sunnyvale, CA 94086
(US). HAFALIA, April [US/US]; 2227 Calle de Primav-
era, Santa Clara, CA 95054 (US). POLICKY, Jennifer,
L. [US/US]; 1511 Jarvis Court, San Jose, CA 95118 (US).(74) Agents: HAMLET-COX, Diana et al.; Incyte Genomics,
Inc., 3160 Porter Drive, Palo Alto, CA 94304 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK,
LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX,
MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL,
TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: EXTRACELLULAR MESSENGERS

(57) Abstract: The invention provides human extracellular messengers (XMES) and polynucleotides which identify and encode XMES. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating or preventing disorders associated with aberrant expression of XMES.



WO 01/94587 A2

EXTRACELLULAR MESSENGERS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of extracellular messengers and
5 to the use of these sequences in the diagnosis, treatment, and prevention of neurological disorders, autoimmune/inflammatory disorders, developmental disorders, endocrine disorders, and cell proliferative disorders including cancer, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of extracellular messengers.

BACKGROUND OF THE INVENTION

Intercellular communication is essential for the growth and survival of multicellular organisms, and in particular, for the function of the endocrine, nervous, and immune systems. In addition, intercellular communication is critical for developmental processes such as tissue construction and organogenesis, in which cell proliferation, cell differentiation, and morphogenesis
15 must be spatially and temporally regulated in a precise and coordinated manner. Cells communicate with one another through the secretion and uptake of diverse types of signaling molecules such as hormones, growth factors, neuropeptides, and cytokines.

Hormones

Hormones are signaling molecules that coordinately regulate basic physiological processes
20 from embryogenesis throughout adulthood. These processes include metabolism, respiration, reproduction, excretion, fetal tissue differentiation and organogenesis, growth and development, homeostasis, and the stress response. Hormonal secretions and the nervous system are tightly integrated and interdependent. Hormones are secreted by endocrine glands, primarily the hypothalamus and pituitary, the thyroid and parathyroid, the pancreas, the adrenal glands, and the ovaries and testes.

25 The secretion of hormones into the circulation is tightly controlled. Hormones are often secreted in diurnal, pulsatile, and cyclic patterns. Hormone secretion is regulated by perturbations in blood biochemistry, by other upstream-acting hormones, by neural impulses, and by negative feedback loops. Blood hormone concentrations are constantly monitored and adjusted to maintain optimal, steady-state levels. Once secreted, hormones act only on those target cells that express specific
30 receptors.

Most disorders of the endocrine system are caused by either hyposecretion or hypersecretion of hormones. Hyposecretion often occurs when a hormone's gland of origin is damaged or otherwise impaired. Hypersecretion often results from the proliferation of tumors derived from hormone-secreting cells. Inappropriate hormone levels may also be caused by defects in regulatory feedback loops or in

the processing of hormone precursors. Endocrine malfunction may also occur when the target cell fails to respond to the hormone.

Hormones can be classified biochemically as polypeptides, steroids, eicosanoids, or amines. Polypeptide hormones, which include diverse hormones such as insulin and growth hormone, vary in size and function and are often synthesized as inactive precursors that are processed intracellularly into mature, active forms. Amine hormones, which include epinephrine and dopamine, are amino acid derivatives that function in neuroendocrine signaling. Steroid hormones, which include the cholesterol-derived hormones estrogen and testosterone, function in sexual development and reproduction. Eicosanoid hormones, which include prostaglandins and prostacyclins, are fatty acid derivatives that function in a variety of processes. Most polypeptide hormones and some amine hormones are soluble in the circulation where they are highly susceptible to proteolytic degradation within seconds after their secretion. Steroid hormones and eicosanoid hormones are insoluble and must be transported in the circulation by carrier proteins. The following discussion will focus primarily on polypeptide hormones.

Hormones secreted by the hypothalamus and pituitary gland play a critical role in endocrine function by coordinately regulating hormonal secretions from other endocrine glands in response to neural signals. Hypothalamic hormones include thyrotropin-releasing hormone, gonadotropin-releasing hormone, somatostatin, growth-hormone releasing factor, corticotropin-releasing hormone, substance P, dopamine, and prolactin-releasing hormone. These hormones directly regulate the secretion of hormones from the anterior lobe of the pituitary. Hormones secreted by the anterior pituitary include adrenocorticotrophic hormone (ACTH), melanocyte-stimulating hormone, somatotrophic hormones such as growth hormone and prolactin, glycoprotein hormones such as thyroid-stimulating hormone, luteinizing hormone (LH), and follicle-stimulating hormone (FSH), β -lipotropin, and β -endorphins. These hormones regulate hormonal secretions from the thyroid, pancreas, and adrenal glands, and act directly on the reproductive organs to stimulate ovulation and spermatogenesis. The posterior pituitary synthesizes and secretes antidiuretic hormone (ADH, vasopressin) and oxytocin.

Disorders of the hypothalamus and pituitary often result from lesions such as primary brain tumors, adenomas, infarction associated with pregnancy, hypophysectomy, aneurysms, vascular malformations, thrombosis, infections, immunological disorders, and complications due to head trauma. Such disorders have profound effects on the function of other endocrine glands. Disorders associated with hypopituitarism include hypogonadism, Sheehan syndrome, diabetes insipidus, Kallman's disease, Hand-Schuller-Christian disease, Letterer-Siwe disease, sarcoidosis, empty sella syndrome, and dwarfism. Disorders associated with hyperpituitarism include acromegaly, gigantism, and syndrome of inappropriate ADH secretion (SIADH), often caused by benign adenomas.

Hormones secreted by the thyroid and parathyroid primarily control metabolic rates and the

regulation of serum calcium levels, respectively. Thyroid hormones include calcitonin, somatostatin, and thyroid hormone. The parathyroid secretes parathyroid hormone. Disorders associated with hypothyroidism include goiter, myxedema, acute thyroiditis associated with bacterial infection, subacute thyroiditis associated with viral infection, autoimmune thyroiditis (Hashimoto's disease), and cretinism. Disorders associated with hyperthyroidism include thyrotoxicosis and its various forms, Grave's disease, pretibial myxedema, toxic multinodular goiter, thyroid carcinoma, and Plummer's disease. Disorders associated with hyperparathyroidism include Conn disease (chronic hypercalcemia) leading to bone resorption and parathyroid hyperplasia.

Hormones secreted by the pancreas regulate blood glucose levels by modulating the rates of carbohydrate, fat, and protein metabolism. Pancreatic hormones include insulin, glucagon, amylin, γ -aminobutyric acid, gastrin, somatostatin, and pancreatic polypeptide. The principal disorder associated with pancreatic dysfunction is diabetes mellitus caused by insufficient insulin activity. Diabetes mellitus is generally classified as either Type I (insulin-dependent, juvenile diabetes) or Type II (non-insulin-dependent, adult diabetes). The treatment of both forms by insulin replacement therapy is well known. Diabetes mellitus often leads to acute complications such as hypoglycemia (insulin shock), coma, diabetic ketoacidosis, lactic acidosis, and chronic complications leading to disorders of the eye, kidney, skin, bone, joint, cardiovascular system, nervous system, and to decreased resistance to infection.

The anatomy, physiology, and diseases related to hormonal function are reviewed in McCance, K. L. and Huether, S. E. (1994) Pathophysiology: The Biological Basis for Disease in Adults and Children, Mosby-Year Book, Inc., St. Louis, MO; Greenspan, F. S. and Baxter, J. D. (1994) Basic and Clinical Endocrinology, Appleton and Lange, East Norwalk, CT.

Growth Factors

Growth factors are secreted proteins that mediate intercellular communication. Unlike hormones, which travel great distances via the circulatory system, most growth factors are primarily local mediators that act on neighboring cells. Most growth factors contain a hydrophobic N-terminal signal peptide sequence which directs the growth factor into the secretory pathway. Most growth factors also undergo post-translational modifications within the secretory pathway. These modifications can include proteolysis, glycosylation, phosphorylation, and intramolecular disulfide bond formation. Once secreted, growth factors bind to specific receptors on the surfaces of neighboring target cells, and the bound receptors trigger intracellular signal transduction pathways. These signal transduction pathways elicit specific cellular responses in the target cells. These responses can include the modulation of gene expression and the stimulation or inhibition of cell division, cell differentiation, and cell motility.

Growth factors fall into at least two broad and overlapping classes. The broadest class

includes the large polypeptide growth factors, which are wide-ranging in their effects. These factors include epidermal growth factor (EGF), fibroblast growth factor (FGF), transforming growth factor- β (TGF- β), insulin-like growth factor (IGF), nerve growth factor (NGF), and platelet-derived growth factor (PDGF), each defining a family of numerous related factors. The large polypeptide growth factors, with the exception of NGF, act as mitogens on diverse cell types to stimulate wound healing, bone synthesis and remodeling, extracellular matrix synthesis, and proliferation of epithelial, epidermal, and connective tissues. Members of the TGF- β , EGF, and FGF families also function as inductive signals in the differentiation of embryonic tissue. NGF functions specifically as a neurotrophic factor, promoting neuronal growth and differentiation.

Another class of growth factors includes the hematopoietic growth factors, which are narrow in their target specificity. These factors stimulate the proliferation and differentiation of blood cells such as B-lymphocytes, T-lymphocytes, erythrocytes, platelets, eosinophils, basophils, neutrophils, macrophages, and their stem cell precursors. These factors include the colony-stimulating factors (G-CSF, M-CSF, GM-CSF, and CSF1-3), erythropoietin, and the cytokines. The cytokines are specialized hematopoietic factors secreted by cells of the immune system and are discussed in detail below.

Growth factors play critical roles in neoplastic transformation of cells in vitro and in tumor progression in vivo. Overexpression of the large polypeptide growth factors promotes the proliferation and transformation of cells in culture. Inappropriate expression of these growth factors by tumor cells in vivo may contribute to tumor vascularization and metastasis. Inappropriate activity of hematopoietic growth factors can result in anemias, leukemias, and lymphomas. Moreover, growth factors are both structurally and functionally related to oncoproteins, the potentially cancer-causing products of proto-oncogenes. Certain FGF and PDGF family members are themselves homologous to oncoproteins, whereas receptors for some members of the EGF, NGF, and FGF families are encoded by proto-oncogenes. Growth factors also affect the transcriptional regulation of both proto-oncogenes and oncosuppressor genes. (Pimentel, E. (1994) Handbook of Growth Factors, CRC Press, Ann Arbor, MI; McKay, I. and Leigh, I., eds. (1993) Growth Factors: A Practical Approach, Oxford University Press, New York, NY; Habenicht, A., ed. (1990) Growth Factors, Differentiation Factors, and Cytokines, Springer-Verlag, New York, NY.)

In addition, some of the large polypeptide growth factors play crucial roles in the induction of the primordial germ layers in the developing embryo. This induction ultimately results in the formation of the embryonic mesoderm, ectoderm, and endoderm which in turn provide the framework for the entire adult body plan. Disruption of this inductive process would be catastrophic to embryonic development.

Small Peptide Factors - Neuropeptides and Vasomediators

Neuropeptides and vasomediators (NP/VM) comprise a family of small peptide factors, typically of 20 amino acids or less. These factors generally function in neuronal excitation and inhibition of vasoconstriction/vasodilation, muscle contraction, and hormonal secretions from the brain and other endocrine tissues. Included in this family are neuropeptides and neuropeptide hormones such as bombesin, neuropeptide Y, neurotensin, neuromedin N, melanocortins, opioids, galanin, somatostatin, tachykinins, urotensin II and related peptides involved in smooth muscle stimulation, vasopressin, vasoactive intestinal peptide, and circulatory system-borne signaling molecules such as angiotensin, complement, calcitonin, endothelins, formyl-methionyl peptides, glucagon, cholecystokinin, gastrin, and many of the peptide hormones discussed above. NP/VMs can transduce signals directly, modulate the activity or release of other neurotransmitters and hormones, and act as catalytic enzymes in signaling cascades. The effects of NP/VMs range from extremely brief to long-lasting. (Reviewed in Martin, C. R. et al. (1985) Endocrine Physiology, Oxford University Press, New York, NY, pp. 57-62.)

15 Cytokines

Cytokines comprise a family of signaling molecules that modulate the immune system and the inflammatory response. Cytokines are usually secreted by leukocytes, or white blood cells, in response to injury or infection. Cytokines function as growth and differentiation factors that act primarily on cells of the immune system such as B- and T-lymphocytes, monocytes, macrophages, and granulocytes. Like other signaling molecules, cytokines bind to specific plasma membrane receptors and trigger intracellular signal transduction pathways which alter gene expression patterns. There is considerable potential for the use of cytokines in the treatment of inflammation and immune system disorders.

Cytokine structure and function have been extensively characterized in vitro. Most cytokines are small polypeptides of about 30 kilodaltons or less. Over 50 cytokines have been identified from human and rodent sources. Examples of cytokine subfamilies include the interferons (IFN- α , - β , and - γ), the interleukins (IL1-IL13), the tumor necrosis factors (TNF- α and - β), and the chemokines. Many cytokines have been produced using recombinant DNA techniques, and the activities of individual cytokines have been determined in vitro. These activities include regulation of leukocyte proliferation, differentiation, and motility.

The activity of an individual cytokine in vitro may not reflect the full scope of that cytokine's activity in vivo. Cytokines are not expressed individually in vivo but are instead expressed in combination with a multitude of other cytokines when the organism is challenged with a stimulus. Together, these cytokines collectively modulate the immune response in a manner appropriate for that particular stimulus. Therefore, the physiological activity of a cytokine is determined by the stimulus

itself and by complex interactive networks among co-expressed cytokines which may demonstrate both synergistic and antagonistic relationships.

Chemokines comprise a cytokine subfamily with over 30 members. (Reviewed in Wells, T. N. C. and Peitsch, M. C. (1997) *J. Leukoc. Biol.* 61:545-550.) Chemokines were initially identified as chemotactic proteins that recruit monocytes and macrophages to sites of inflammation. Recent evidence indicates that chemokines may also play key roles in hematopoiesis and HIV-1 infection. Chemokines are small proteins which range from about 6-15 kilodaltons in molecular weight. Chemokines are further classified as C, CC, CXC, or CX₃C based on the number and position of critical cysteine residues. The CC chemokines, for example, each contain a conserved motif consisting of two consecutive cysteines followed by two additional cysteines which occur downstream at 24- and 16-residue intervals, respectively (ExpASY PROSITE database, documents PS00472 and PDOC00434). The presence and spacing of these four cysteine residues are highly conserved, whereas the intervening residues diverge significantly. However, a conserved tyrosine located about 15 residues downstream of the cysteine doublet seems to be important for chemotactic activity. Most of the human genes encoding CC chemokines are clustered on chromosome 17, although there are a few examples of CC chemokine genes that map elsewhere. Other chemokines include lymphotactin (C chemokine); macrophage chemotactic and activating factor (MCAF/MCP-1; CC chemokine); platelet factor 4 and IL-8 (CXC chemokines); and fractalkine and neurotractin (CX₃C chemokines). (Reviewed in Luster, A. D. (1998) *N. Engl. J. Med.* 338:436-445.)

The discovery of new extracellular messengers and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of neurological disorders, autoimmune/inflammatory disorders, developmental disorders, endocrine disorders, and cell proliferative disorders including cancer, and in the assessment of the effects of exogenous compounds on the expression of nucleic acid and amino acid sequences of extracellular messengers.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, extracellular messengers, referred to collectively as "XMES" and individually as "XMES-1," "XMES-2," "XMES-3," "XMES-4," "XMES-5," "XMES-6," "XMES-7," "XMES-8," and "XMES-9." In one aspect, the invention provides an isolated polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid

sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-9.

5 The invention further provides an isolated polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. In one alternative, the polynucleotide encodes a polypeptide selected from the group consisting of SEQ ID NO:1-9. In another alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:10-18.

15 Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

25 The invention also provides a method for producing a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

30 Additionally, the invention provides an isolated antibody which specifically binds to a

polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9.

The invention further provides an isolated polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide or fragments thereof, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide selected from the group consisting of a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, c) a polynucleotide complementary to the polynucleotide of a), d) a polynucleotide complementary to the polynucleotide of b), and e) an RNA equivalent of a)-d). The method comprises a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and b)

detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.

The invention further provides a composition comprising an effective amount of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and a pharmaceutically acceptable excipient. In one embodiment, the composition comprises an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional XMES, comprising administering to a patient in need of such treatment the composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional XMES, comprising administering to a patient in need of such treatment the composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. The method comprises a) exposing a sample comprising the

polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional XMES, comprising
5 administering to a patient in need of such treatment the composition.

The invention further provides a method of screening for a compound that specifically binds to a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the
10 group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. The method comprises a) combining the polypeptide with at least one test compound under suitable conditions, and b) detecting binding of the polypeptide to the test compound, thereby identifying a
15 compound that specifically binds to the polypeptide.

The invention further provides a method of screening for a compound that modulates the activity of a polypeptide selected from the group consisting of a) a polypeptide comprising an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence
20 selected from the group consisting of SEQ ID NO:1-9, c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9. The method comprises a) combining the polypeptide with at least one test compound under conditions permissive for the activity of the polypeptide, b) assessing the
25 activity of the polypeptide in the presence of the test compound, and c) comparing the activity of the polypeptide in the presence of the test compound with the activity of the polypeptide in the absence of the test compound, wherein a change in the activity of the polypeptide in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide.

The invention further provides a method for screening a compound for effectiveness in
30 altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:10-18, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

The invention further provides a method for assessing toxicity of a test compound, said

method comprising a) treating a biological sample containing nucleic acids with the test compound;
b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20
contiguous nucleotides of a polynucleotide selected from the group consisting of i) a polynucleotide
comprising a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, ii) a
5 naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a
polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, iii) a
polynucleotide having a sequence complementary to i), iv) a polynucleotide complementary to the
polynucleotide of ii), and v) an RNA equivalent of i)-iv). Hybridization occurs under conditions
whereby a specific hybridization complex is formed between said probe and a target polynucleotide in
10 the biological sample, said target polynucleotide selected from the group consisting of i) a
polynucleotide comprising a polynucleotide sequence selected from the group consisting of SEQ ID
NO:10-18, ii) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90%
identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18, iii) a
polynucleotide complementary to the polynucleotide of i), iv) a polynucleotide complementary to the
15 polynucleotide of ii), and v) an RNA equivalent of i)-iv). Alternatively, the target polynucleotide
comprises a fragment of a polynucleotide sequence selected from the group consisting of i)-v) above;
c) quantifying the amount of hybridization complex; and d) comparing the amount of hybridization
complex in the treated biological sample with the amount of hybridization complex in an untreated
biological sample, wherein a difference in the amount of hybridization complex in the treated
20 biological sample is indicative of toxicity of the test compound.

BRIEF DESCRIPTION OF THE TABLES

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide
sequences of the present invention.

25 Table 2 shows the GenBank identification number and annotation of the nearest GenBank
homolog for polypeptides of the invention. The probability score for the match between each
polypeptide and its GenBank homolog is also shown.

Table 3 shows structural features of polypeptide sequences of the invention, including predicted
motifs and domains, along with the methods, algorithms, and searchable databases used for analysis of
30 the polypeptides.

Table 4 lists the cDNA and/or genomic DNA fragments which were used to assemble
polynucleotide sequences of the invention, along with selected fragments of the polynucleotide
sequences.

Table 5 shows the representative cDNA library for polynucleotides of the invention.

Table 6 provides an appendix which describes the tissues and vectors used for construction of the cDNA libraries shown in Table 5.

Table 7 shows the tools, programs, and algorithms used to analyze the polynucleotides and polypeptides of the invention, along with applicable descriptions, references, and threshold parameters.

5

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the invention is not entitled to antedate such disclosure by virtue of prior invention.

25 DEFINITIONS

"XMES" refers to the amino acid sequences of substantially purified XMES obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

The term "agonist" refers to a molecule which intensifies or mimics the biological activity of XMES. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of XMES either by directly interacting with XMES or by acting on components of the biological pathway in which XMES participates.

An "allelic variant" is an alternative form of the gene encoding XMES. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in

polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

“Altered” nucleic acid sequences encoding XMES include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as XMES or a polypeptide with at least one functional characteristic of XMES. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding XMES, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding XMES. The encoded protein may also be “altered,” and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent XMES. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of XMES is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms “amino acid” and “amino acid sequence” refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where “amino acid sequence” is recited to refer to a sequence of a naturally occurring protein molecule, “amino acid sequence” and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

“Amplification” relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term “antagonist” refers to a molecule which inhibits or attenuates the biological activity of XMES. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of XMES either by directly interacting with XMES or by acting on components of the biological pathway in which XMES participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind XMES polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition capable of base-pairing with the "sense" (coding) strand of a specific nucleic acid sequence. Antisense compositions may include DNA; RNA; peptide nucleic acid (PNA); oligonucleotides having modified backbone linkages such as phosphorothioates, methylphosphonates, or benzylphosphonates; oligonucleotides having modified sugar groups such as 2'-methoxyethyl sugars or 2'-methoxyethoxy sugars; or oligonucleotides having modified bases such as 5-methyl cytosine, 2'-deoxyuracil, or 7-deaza-2'-deoxyguanosine. Antisense molecules may be produced by any method including chemical synthesis or transcription. Once introduced into a cell, the complementary antisense molecule base-pairs with a naturally occurring nucleic acid sequence produced by the cell to form duplexes which block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand of a reference DNA molecule.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" or "immunogenic" refers to the capability of the natural, recombinant, or synthetic XMES, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

"Complementary" describes the relationship between two single-stranded nucleic acid sequences that anneal by base-pairing. For example, 5'-AGT-3' pairs with its complement, 3'-TCA-5'.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a

given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding XMES or fragments of XMES may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been subjected to repeated DNA sequence analysis to resolve uncalled bases, extended using the XL-PCR kit (Applied Biosystems, Foster City CA) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from one or more overlapping cDNA, EST, or genomic DNA fragments using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI) or Phrap (University of Washington, Seattle WA). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that are predicted to least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
	Thr	Ser, Val
	Trp	Phe, Tyr
	Tyr	His, Phe, Trp
	Val	Ile, Leu, Thr

Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide

backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

5 A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

The term "derivative" refers to a chemically modified polynucleotide or polypeptide. Chemical modifications of a polynucleotide can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified
10 by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

A "detectable label" refers to a reporter molecule or enzyme that is capable of generating a measurable signal and is covalently or noncovalently joined to a polynucleotide or polypeptide.

"Differential expression" refers to increased or upregulated; or decreased, downregulated, or
15 absent gene or protein expression, determined by comparing at least two different samples. Such comparisons may be carried out between, for example, a treated and an untreated sample, or a diseased and a normal sample.

A "fragment" is a unique portion of XMES or the polynucleotide encoding XMES which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up
20 to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 16, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For
25 example, a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50%) of a polypeptide as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

30 A fragment of SEQ ID NO:10-18 comprises a region of unique polynucleotide sequence that specifically identifies SEQ ID NO:10-18, for example, as distinct from any other sequence in the genome from which the fragment was obtained. A fragment of SEQ ID NO:10-18 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:10-18 from related polynucleotide sequences. The precise length of a fragment of SEQ

ID NO:10-18 and the region of SEQ ID NO:10-18 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A fragment of SEQ ID NO:1-9 is encoded by a fragment of SEQ ID NO:10-18. A fragment of SEQ ID NO:1-9 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-9. For example, a fragment of SEQ ID NO:1-9 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-9. The precise length of a fragment of SEQ ID NO:1-9 and the region of SEQ ID NO:1-9 to which the fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

A "full length" polynucleotide sequence is one containing at least a translation initiation codon (e.g., methionine) followed by an open reading frame and a translation termination codon. A "full length" polynucleotide sequence encodes a "full length" polypeptide sequence.

"Homology" refers to sequence similarity or, interchangeably, sequence identity, between two or more polynucleotide sequences or two or more polypeptide sequences.

The terms "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191.

For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polynucleotide sequences.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD, and on the Internet at <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other

polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Reward for match: 1

10 *Penalty for mismatch: -2*

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 11

15 *Filter: on*

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the charge and hydrophobicity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment

program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.12 (April-21-2000) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size and which contain all of the elements required for chromosome replication, segregation and maintenance.

The term "humanized antibody" refers to an antibody molecule in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of complementarity. Specific hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for

annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml sheared, denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Such wash temperatures are typically selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour. Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, sheared and denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of

various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect cellular and systemic defense systems.

An "immunogenic fragment" is a polypeptide or oligopeptide fragment of XMES which is capable of eliciting an immune response when introduced into a living organism, for example, a mammal. The term "immunogenic fragment" also includes any polypeptide or oligopeptide fragment of XMES which is useful in any of the antibody production methods disclosed herein or known in the art.

The term "microarray" refers to an arrangement of a plurality of polynucleotides, polypeptides, or other chemical compounds on a substrate.

The terms "element" and "array element" refer to a polynucleotide, polypeptide, or other chemical compound having a unique and defined position on a microarray.

The term "modulate" refers to a change in the activity of XMES. For example, modulation may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of XMES.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with a second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Operably linked DNA sequences may be in close proximity or contiguous and, where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation, and may be pegylated to extend their lifespan in the cell.

"Post-translational modification" of an XMES may involve lipidation, glycosylation, phosphorylation, acetylation, racemization, proteolytic cleavage, and other modifications known in the art. These processes may occur synthetically or biochemically. Biochemical modifications will vary by cell type depending on the enzymatic milieu of XMES.

"Probe" refers to nucleic acid sequences encoding XMES, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical

labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes. "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

Methods for preparing and using probes and primers are described in the references, for example Sambrook, J. et al. (1989) Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel, F.M. et al. (1987) Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis, M. et al. (1990) PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that

hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing
5 primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence that is made by an artificial combination of two or more otherwise separated segments of sequence.
10 This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence.
15 Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

A "regulatory element" refers to a nucleic acid sequence usually derived from untranslated
20 regions of a gene and includes enhancers, promoters, introns, and 5' and 3' untranslated regions (UTRs). Regulatory elements interact with host or viral proteins which control transcription, translation, or RNA stability.

"Reporter molecules" are chemical or biochemical moieties used for labeling a nucleic acid, amino acid, or antibody. Reporter molecules include radionuclides; enzymes; fluorescent,
25 chemiluminescent, or chromogenic agents; substrates; cofactors; inhibitors; magnetic particles; and other moieties known in the art.

An "RNA equivalent," in reference to a DNA sequence, is composed of the same linear sequence of nucleotides as the reference DNA sequence with the exception that all occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose
30 instead of deoxyribose.

The term "sample" is used in its broadest sense. A sample suspected of containing XMES, nucleic acids encoding XMES, or fragments thereof may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For
5 example, if an antibody is specific for epitope "A," the presence of a polypeptide comprising the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free,
10 preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acid residues or nucleotides by different amino acid residues or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters,
15 chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

A "transcript image" refers to the collective pattern of gene expression by a particular cell type or tissue under given conditions at a given time.

20 "Transformation" describes a process by which exogenous DNA is introduced into a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected based on the type of host cell being transformed and may include, but is not limited to, bacteriophage or viral infection,
25 electroporation, heat shock, lipofection, and particle bombardment. The term "transformed cells" includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently transformed cells which express the inserted DNA or RNA for limited periods of time.

A "transgenic organism," as used herein, is any organism, including but not limited to
30 animals and plants, in which one or more of the cells of the organism contains heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in

vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. The transgenic organisms contemplated in accordance with the present invention include bacteria, cyanobacteria, fungi, plants and animals. The isolated DNA of the present invention can be introduced into the host by methods known in the art, for example infection, transfection, transformation or transconjugation. Techniques for transferring the DNA of the present invention into such organisms are widely known and provided in references such as Sambrook et al. (1989), supra.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternative splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides will generally have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 91%, at least 92%, at least 93%, at least 94%, at least 95%, at least 96%, at least 97%, at least 98%, or at least 99% or greater sequence identity over a certain defined length of one of the polypeptides.

THE INVENTION

The invention is based on the discovery of new human extracellular messengers (XMES), the

polynucleotides encoding XMES, and the use of these compositions for the diagnosis, treatment, or prevention of neurological disorders, autoimmune/inflammatory disorders, developmental disorders, endocrine disorders, and cell proliferative disorders including cancer.

Table 1 summarizes the nomenclature for the full length polynucleotide and polypeptide sequences of the invention. Each polynucleotide and its corresponding polypeptide are correlated to a single Incyte project identification number (Incyte Project ID). Each polypeptide sequence is denoted by both a polypeptide sequence identification number (Polypeptide SEQ ID NO:) and an Incyte polypeptide sequence number (Incyte Polypeptide ID) as shown. Each polynucleotide sequence is denoted by both a polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and an Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) as shown.

Table 2 shows sequences with homology to the polypeptides of the invention as identified by BLAST analysis against the GenBank protein (genpept) database. Columns 1 and 2 show the polypeptide sequence identification number (Polypeptide SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for polypeptides of the invention. Column 3 shows the GenBank identification number (Genbank ID NO:) of the nearest GenBank homolog. Column 4 shows the probability score for the match between each polypeptide and its GenBank homolog. Column 5 shows the annotation of the GenBank homolog along with relevant citations where applicable, all of which are expressly incorporated by reference herein.

Table 3 shows various structural features of the polypeptides of the invention. Columns 1 and 2 show the polypeptide sequence identification number (SEQ ID NO:) and the corresponding Incyte polypeptide sequence number (Incyte Polypeptide ID) for each polypeptide of the invention. Column 3 shows the number of amino acid residues in each polypeptide. Column 4 shows potential phosphorylation sites, and column 5 shows potential glycosylation sites, as determined by the MOTIFS program of the GCG sequence analysis software package (Genetics Computer Group, Madison WI). Column 6 shows amino acid residues comprising signature sequences, domains, and motifs. Column 7 shows analytical methods for protein structure/function analysis and in some cases, searchable databases to which the analytical methods were applied.

Together, Tables 2 and 3 summarize the properties of polypeptides of the invention, and these properties establish that the claimed polypeptides are extracellular messengers. For example, SEQ ID NO:2 is 96% identical, over 1210 amino acids, to rat neurexin III-alpha (GenBank ID g394600) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is 0.0, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. Based on this analysis, SEQ ID NO:2 is a human neurexin. In an alternative example, SEQ ID NO:1 is 39% identical over 240 amino acids to mac25 (g3721617) (a protein which

binds the class of proteins known as insulin-like growth factors) as determined by the Basic Local Alignment Search Tool (BLAST). The probability score is $3.3\text{e-}42$ (see Table 2). In an additional alternative example, SEQ ID NO:3 is 39% identical, to human fibroblast growth factor 19 (GenBank ID g4514718) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The

5 BLAST probability score is $1.5\text{e-}22$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID NO:3 also contains a fibroblast growth factor domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from

10 BLIMPS and PROFILESCAN analyses provide further corroborative evidence that SEQ ID NO:3 is a fibroblast growth factor. In another alternative example, SEQ ID NO:6 is 81% identical to Bos taurus cartilage-derived morphogenetic protein (GenBank ID g632490) as determined by the Basic Local Alignment Search Tool (BLAST). (See Table 2.) The BLAST probability score is $7.5\text{e-}185$, which indicates the probability of obtaining the observed polypeptide sequence alignment by chance. SEQ ID

15 NO:6 also contains a TGF-beta propeptide domain and a TGF-beta-like domain as determined by searching for statistically significant matches in the hidden Markov model (HMM)-based PFAM database of conserved protein family domains. (See Table 3.) Data from BLIMPS analysis provides further corroborative evidence that SEQ ID NO:6 is a TGF-beta family protein. SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:7, SEQ ID NO:8 and SEQ ID NO:9 were analyzed and annotated in a similar manner. The algorithms and parameters for the analysis of SEQ ID NO:1-9 are described in Table 7.

20 As shown in Table 4, the full length polynucleotide sequences of the present invention were assembled using cDNA sequences or coding (exon) sequences derived from genomic DNA, or any combination of these two types of sequences. Columns 1 and 2 list the polynucleotide sequence identification number (Polynucleotide SEQ ID NO:) and the corresponding Incyte polynucleotide consensus sequence number (Incyte Polynucleotide ID) for each polynucleotide of the invention.

25 Column 3 shows the length of each polynucleotide sequence in basepairs. Column 4 lists fragments of the polynucleotide sequences which are useful, for example, in hybridization or amplification technologies that identify SEQ ID NO:10-18 or that distinguish between SEQ ID NO:10-18 and related polynucleotide sequences. Column 5 shows identification numbers corresponding to cDNA sequences, coding sequences (exons) predicted from genomic DNA, and/or sequence assemblages

30 comprised of both cDNA and genomic DNA. These sequences were used to assemble the full length polynucleotide sequences of the invention. Columns 6 and 7 of Table 4 show the nucleotide start (5') and stop (3') positions of the cDNA and/or genomic sequences in column 5 relative to their respective full length sequences.

The identification numbers in Column 5 of Table 4 may refer specifically, for example, to

Incyte cDNAs along with their corresponding cDNA libraries. Incyte cDNAs for which cDNA libraries are not indicated were derived from pooled cDNA libraries (e.g., 70656442V1). Alternatively, the identification numbers in column 5 may refer to GenBank cDNAs or ESTs (e.g., g3153915) which contributed to the assembly of the full length polynucleotide sequences. Alternatively, the identification numbers in column 5 may refer to coding regions predicted by Genscan analysis of genomic DNA. For example, GNN.g7768040_000033_002 is the identification number of a Genscan-predicted coding sequence, with g7768040 being the GenBank identification number of the sequence to which Genscan was applied. The Genscan-predicted coding sequences may have been edited prior to assembly. (See Example IV.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon stitching" algorithm. (See Example V.) Alternatively, the identification numbers in column 5 may refer to assemblages of both cDNA and Genscan-predicted exons brought together by an "exon-stretching" algorithm. For example, FL5378618_g4416547_g4091819 is the identification number of a "stretched" sequence, with 5378618 being the Incyte project identification number, g4416547 being the GenBank identification number of the human genomic sequence to which the "exon-stretching" algorithm was applied, and g4091819 being the GenBank identification number of the nearest GenBank protein homolog. (See Example V.) In some cases, Incyte cDNA coverage redundant with the sequence coverage shown in column 5 was obtained to confirm the final consensus polynucleotide sequence, but the relevant Incyte cDNA identification numbers are not shown.

Table 5 shows the representative cDNA libraries for those full length polynucleotide sequences which were assembled using Incyte cDNA sequences. The representative cDNA library is the Incyte cDNA library which is most frequently represented by the Incyte cDNA sequences which were used to assemble and confirm the above polynucleotide sequences. The tissues and vectors which were used to construct the cDNA libraries shown in Table 5 are described in Table 6.

The invention also encompasses XMES variants. A preferred XMES variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the XMES amino acid sequence, and which contains at least one functional or structural characteristic of XMES.

The invention also encompasses polynucleotides which encode XMES. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:10-18, which encodes XMES. The polynucleotide sequences of SEQ ID NO:10-18, as presented in the Sequence Listing, embrace the equivalent RNA sequences, wherein occurrences of the nitrogenous base thymine are replaced with uracil, and the sugar backbone is composed of ribose instead of deoxyribose.

The invention also encompasses a variant of a polynucleotide sequence encoding XMES. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding XMES. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:10-18 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:10-18. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of XMES.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding XMES, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring XMES, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode XMES and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring XMES under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding XMES or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding XMES and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode XMES and XMES derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding XMES or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:10-18 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and

S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the
5 embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of
DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Applied
Biosystems), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or
combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE
amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is
10 automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV),
PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler
(Applied Biosystems). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing
system (Applied Biosystems), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics,
Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a
15 variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols
in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular
Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding XMES may be extended utilizing a partial nucleotide
sequence and employing various PCR-based methods known in the art to detect upstream sequences,
20 such as promoters and regulatory elements. For example, one method which may be employed,
restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic
DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.)
Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown
sequence from a circularized template. The template is derived from restriction fragments comprising a
25 known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids*
Res. 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent
to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al.
(1991) *PCR Methods Applic.* 1:111-119.) In this method, multiple restriction enzyme digestions and
ligations may be used to insert an engineered double-stranded sequence into a region of unknown
30 sequence before performing PCR. Other methods which may be used to retrieve unknown sequences
are known in the art. (See, e.g., Parker, J.D. et al. (1991) *Nucleic Acids Res.* 19:3055-3060).
Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo
Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in
finding intron/exon junctions. For all PCR-based methods, primers may be designed using

commercially available software, such as OLIGO 4.06 primer analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

5 When screening for full length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

10 Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate
15 software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Applied Biosystems), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be present in limited amounts in a particular sample.

 In another embodiment of the invention, polynucleotide sequences or fragments thereof which
20 encode XMES may be cloned in recombinant DNA molecules that direct expression of XMES, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a functionally equivalent amino acid sequence may be produced and used to express XMES.

 The nucleotide sequences of the present invention can be engineered using methods generally
25 known in the art in order to alter XMES-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites,
30 alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

 The nucleotides of the present invention may be subjected to DNA shuffling techniques such as MOLECULARBREEDING (Maxygen Inc., Santa Clara CA; described in U.S. Patent Number 5,837,458; Chang, C.-C. et al. (1999) Nat. Biotechnol. 17:793-797; Christians, F.C. et al. (1999) Nat. Biotechnol. 17:259-264; and Crameri, A. et al. (1996) Nat. Biotechnol. 14:315-319) to alter or

improve the biological properties of XMES, such as its biological or enzymatic activity or its ability to bind to other molecules or compounds. DNA shuffling is a process by which a library of gene variants is produced using PCR-mediated recombination of gene fragments. The library is then subjected to selection or screening procedures that identify those gene variants with the desired properties. These preferred variants may then be pooled and further subjected to recursive rounds of DNA shuffling and selection/screening. Thus, genetic diversity is created through "artificial" breeding and rapid molecular evolution. For example, fragments of a single gene containing random point mutations may be recombined, screened, and then reshuffled until the desired properties are optimized. Alternatively, fragments of a given gene may be recombined with fragments of homologous genes in the same gene family, either from the same or different species, thereby maximizing the genetic diversity of multiple naturally occurring genes in a directed and controllable manner.

In another embodiment, sequences encoding XMES may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) Nucleic Acids Symp. Ser. 7:215-223; and Horn, T. et al. (1980) Nucleic Acids Symp. Ser. 7:225-232.) Alternatively, XMES itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solution-phase or solid-phase techniques. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY, pp. 55-60; and Roberge, J.Y. et al. (1995) Science 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Applied Biosystems). Additionally, the amino acid sequence of XMES, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide or a polypeptide having a sequence of a naturally occurring polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) Methods Enzymol. 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, supra, pp. 28-53.)

In order to express a biologically active XMES, the nucleotide sequences encoding XMES or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding XMES. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding XMES. Such signals

include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding XMES and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous
5 translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

10 Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding XMES and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995)
15 Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

A variety of expression vector/host systems may be utilized to contain and express sequences encoding XMES. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus);
20 plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or animal cell systems. (See, e.g., Sambrook, supra; Ausubel, supra; Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509; Engelhard, E.K. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3224-3227; Sandig, V. et al. (1996) *Hum. Gene Ther.* 7:1937-1945; Takamatsu, N. (1987) *EMBO J.* 6:307-311; The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New
25 York NY, pp. 191-196; Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659; and Harrington, J.J. et al. (1997) *Nat. Genet.* 15:345-355.) Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. (See, e.g., Di
30 Nicola, M. et al. (1998) *Cancer Gen. Ther.* 5(6):350-356; Yu, M. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90(13):6340-6344; Buller, R.M. et al. (1985) *Nature* 317(6040):813-815; McGregor, D.P. et al. (1994) *Mol. Immunol.* 31(3):219-226; and Verma, I.M. and N. Somia (1997) *Nature* 389:239-242.) The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending

upon the use intended for polynucleotide sequences encoding XMES. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding XMES can be achieved using a multifunctional *E. coli* vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding XMES into the vector's multiple cloning site
5 disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for *in vitro* transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) *J. Biol. Chem.* 264:5503-5509.) When large quantities of XMES are needed, e.g. for the production of antibodies,
10 vectors which direct high level expression of XMES may be used. For example, vectors containing the strong, inducible SP6 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of XMES. A number of vectors containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast *Saccharomyces cerevisiae* or *Pichia pastoris*. In addition, such
15 vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, *supra*; Bitter, G.A. et al. (1987) *Methods Enzymol.* 153:516-544; and Scorer, C.A. et al. (1994) *Bio/Technology* 12:181-184.)

Plant systems may also be used for expression of XMES. Transcription of sequences encoding
20 XMES may be driven by viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) *EMBO J.* 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) *EMBO J.* 3:1671-1680; Broglie, R. et al. (1984) *Science* 224:838-843; and Winter, J. et al. (1991) *Results Probl. Cell Differ.* 17:85-105.) These constructs can
25 be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., *The McGraw Hill Yearbook of Science and Technology* (1992) McGraw Hill, New York NY, pp. 191-196.)

In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding XMES may be ligated into an
30 adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses XMES in host cells. (See, e.g., Logan, J. and T. Shenk (1984) *Proc. Natl. Acad. Sci. USA* 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-

based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of XMES in cell lines is preferred. For example, sequences encoding XMES can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk*⁻ and *apr*⁻ cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding XMES is inserted within a marker gene sequence, transformed cells containing sequences encoding XMES can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding XMES under the control of a single

promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding XMES and that express XMES may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR
5 amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of XMES using either specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include
10 enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on XMES is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See, e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E.
15 et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding XMES include
20 oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding XMES, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of
25 commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding XMES may be cultured under
30 conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode XMES may be designed to contain signal sequences which direct secretion of XMES through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding XMES may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric XMES protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of XMES activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the XMES encoding sequence and the heterologous protein sequence, so that XMES may be cleaved away from the heterologous moiety following purification. Methods for fusion protein expression and purification are discussed in Ausubel (1995, *supra*, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled XMES may be achieved *in vitro* using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

XMES of the present invention or fragments thereof may be used to screen for compounds that specifically bind to XMES. At least one and up to a plurality of test compounds may be screened for specific binding to XMES. Examples of test compounds include antibodies, oligonucleotides,

proteins (e.g., receptors), or small molecules.

In one embodiment, the compound thus identified is closely related to the natural ligand of XMES, e.g., a ligand or fragment thereof, a natural substrate, a structural or functional mimetic, or a natural binding partner. (See, e.g., Coligan, J.E. et al. (1991) Current Protocols in Immunology 1(2): Chapter 5.) Similarly, the compound can be closely related to the natural receptor to which XMES binds, or to at least a fragment of the receptor, e.g., the ligand binding site. In either case, the compound can be rationally designed using known techniques. In one embodiment, screening for these compounds involves producing appropriate cells which express XMES, either as a secreted protein or on the cell membrane. Preferred cells include cells from mammals, yeast, Drosophila, or E. coli. Cells expressing XMES or cell membrane fractions which contain XMES are then contacted with a test compound and binding, stimulation, or inhibition of activity of either XMES or the compound is analyzed.

An assay may simply test binding of a test compound to the polypeptide, wherein binding is detected by a fluorophore, radioisotope, enzyme conjugate, or other detectable label. For example, the assay may comprise the steps of combining at least one test compound with XMES, either in solution or affixed to a solid support, and detecting the binding of XMES to the compound. Alternatively, the assay may detect or measure binding of a test compound in the presence of a labeled competitor. Additionally, the assay may be carried out using cell-free preparations, chemical libraries, or natural product mixtures, and the test compound(s) may be free in solution or affixed to a solid support.

XMES of the present invention or fragments thereof may be used to screen for compounds that modulate the activity of XMES. Such compounds may include agonists, antagonists, or partial or inverse agonists. In one embodiment, an assay is performed under conditions permissive for XMES activity, wherein XMES is combined with at least one test compound, and the activity of XMES in the presence of a test compound is compared with the activity of XMES in the absence of the test compound. A change in the activity of XMES in the presence of the test compound is indicative of a compound that modulates the activity of XMES. Alternatively, a test compound is combined with an in vitro or cell-free system comprising XMES under conditions suitable for XMES activity, and the assay is performed. In either of these assays, a test compound which modulates the activity of XMES may do so indirectly and need not come in direct contact with the test compound. At least one and up to a plurality of test compounds may be screened.

In another embodiment, polynucleotides encoding XMES or their mammalian homologs may be "knocked out" in an animal model system using homologous recombination in embryonic stem (ES) cells. Such techniques are well known in the art and are useful for the generation of animal models of

human disease. (See, e.g., U.S. Patent Number 5,175,383 and U.S. Patent Number 5,767,337.) For example, mouse ES cells, such as the mouse 129/SvJ cell line, are derived from the early mouse embryo and grown in culture. The ES cells are transformed with a vector containing the gene of interest disrupted by a marker gene, e.g., the neomycin phosphotransferase gene (neo; Capecchi, M.R. (1989) Science 244:1288-1292). The vector integrates into the corresponding region of the host genome by homologous recombination. Alternatively, homologous recombination takes place using the Cre-loxP system to knockout a gene of interest in a tissue- or developmental stage-specific manner (Marth, J.D. (1996) Clin. Invest. 97:1999-2002; Wagner, K.U. et al. (1997) Nucleic Acids Res. 25:4323-4330). Transformed ES cells are identified and microinjected into mouse cell blastocysts such as those from the C57BL/6 mouse strain. The blastocysts are surgically transferred to pseudopregnant dams, and the resulting chimeric progeny are genotyped and bred to produce heterozygous or homozygous strains. Transgenic animals thus generated may be tested with potential therapeutic or toxic agents.

Polynucleotides encoding XMES may also be manipulated in vitro in ES cells derived from human blastocysts. Human ES cells have the potential to differentiate into at least eight separate cell lineages including endoderm, mesoderm, and ectodermal cell types. These cell lineages differentiate into, for example, neural cells, hematopoietic lineages, and cardiomyocytes (Thomson, J.A. et al. (1998) Science 282:1145-1147).

Polynucleotides encoding XMES can also be used to create "knockin" humanized animals (pigs) or transgenic animals (mice or rats) to model human disease. With knockin technology, a region of a polynucleotide encoding XMES is injected into animal ES cells, and the injected sequence integrates into the animal cell genome. Transformed cells are injected into blastulae, and the blastulae are implanted as described above. Transgenic progeny or inbred lines are studied and treated with potential pharmaceutical agents to obtain information on treatment of a human disease. Alternatively, a mammal inbred to overexpress XMES, e.g., by secreting XMES in its milk, may also serve as a convenient source of that protein (Janne, J. et al. (1998) Biotechnol. Annu. Rev. 4:55-74).

THERAPEUTICS

Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of XMES and extracellular messengers. In addition, the expression of XMES is closely associated with cartilage and neurological tissues and with tumor tissues from brain, breast, liver, and prostate. Therefore, XMES appears to play a role in neurological disorders, autoimmune/inflammatory disorders, developmental disorders, endocrine disorders, and cell proliferative disorders including cancer. In the treatment of disorders associated with increased XMES expression or activity, it is desirable to decrease the expression or activity of XMES. In the treatment of disorders associated with decreased XMES expression or activity, it is desirable to increase the

expression or activity of XMES.

Therefore, in one embodiment, XMES or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMES. Examples of such disorders include, but are not limited to, a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a developmental disorder such as

renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss; an endocrine disorder such as a disorder of the hypothalamus and/or pituitary resulting from lesions such as a primary brain tumor, adenoma, infarction associated with pregnancy, hypophysectomy, aneurysm, vascular malformation, thrombosis, infection, immunological disorder, and complication due to head trauma; a disorder associated with hypopituitarism including hypogonadism, Sheehan syndrome, diabetes insipidus, Kallman's disease, Hand-Schuller-Christian disease, Letterer-Siwe disease, sarcoidosis, empty sella syndrome, and dwarfism; a disorder associated with hyperpituitarism including acromegaly, gigantism, and syndrome of inappropriate antidiuretic hormone (ADH) secretion (SIADH) often caused by benign adenoma; a disorder associated with hypothyroidism including goiter, myxedema, acute thyroiditis associated with bacterial infection, subacute thyroiditis associated with viral infection, autoimmune thyroiditis (Hashimoto's disease), and cretinism; a disorder associated with hyperthyroidism including thyrotoxicosis and its various forms, Grave's disease, pretibial myxedema, toxic multinodular goiter, thyroid carcinoma, and Plummer's disease; a disorder associated with hyperparathyroidism including Conn disease (chronic hypercalcemia); a pancreatic disorder such as Type I or Type II diabetes mellitus and associated complications; a disorder associated with the adrenals such as hyperplasia, carcinoma, or adenoma of the adrenal cortex, hypertension associated with alkalosis, amyloidosis, hypokalemia, Cushing's disease, Liddle's syndrome, and Arnold-Healy-Gordon syndrome, pheochromocytoma tumors, and Addison's disease; a disorder associated with gonadal steroid hormones such as: in women, abnormal prolactin production, infertility, endometriosis, perturbation of the menstrual cycle, polycystic ovarian disease, hyperprolactinemia, isolated gonadotropin deficiency, amenorrhea, galactorrhea, hermaphroditism, hirsutism and virilization, breast cancer, and, in post-menopausal women, osteoporosis; and, in men, Leydig cell deficiency, male climacteric phase, and germinal cell aplasia, a hypergonadal disorder associated with Leydig cell tumors, androgen resistance associated with absence of androgen receptors, 5 α -reductase deficiency, 21-hydroxylase deficiency, and gynecomastia, and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma,

teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

In another embodiment, a vector capable of expressing XMES or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMES including, but not limited to, those described above.

In a further embodiment, a composition comprising a substantially purified XMES in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMES including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of XMES may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of XMES including, but not limited to, those listed above.

In a further embodiment, an antagonist of XMES may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of XMES. Examples of such disorders include, but are not limited to, those neurological disorders, autoimmune/inflammatory disorders, developmental disorders, endocrine disorders, and cell proliferative disorders including cancer described above. In one aspect, an antibody which specifically binds XMES may be used directly as an antagonist or indirectly as a targeting or delivery mechanism for bringing a pharmaceutical agent to cells or tissues which express XMES.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding XMES may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of XMES including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of XMES may be produced using methods which are generally known in the art. In particular, purified XMES may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind XMES. Antibodies to XMES may also be generated using methods that are well known in the art. Such antibodies may include, but are not

limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans,
5 and others may be immunized by injection with XMES or with any fragment or oligopeptide thereof which has immunogenic properties. Depending on the host species, various adjuvants may be used to increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG
10 (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to XMES have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein. Short stretches of XMES amino
15 acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to XMES may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma
20 technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate
25 antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce XMES-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be
30 generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA 86:3833-3837; Winter,

G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for XMES may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D. et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between XMES and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering XMES epitopes is generally used, but a competitive binding assay may also be employed (Pound, *supra*).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for XMES. Affinity is expressed as an association constant, K_a, which is defined as the molar concentration of XMES-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple XMES epitopes, represents the average affinity, or avidity, of the antibodies for XMES. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular XMES epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10⁹ to 10¹² L/mole are preferred for use in immunoassays in which the XMES-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10⁶ to 10⁷ L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of XMES, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington DC; Liddell, J.E. and A. Cryer (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of XMES-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, *supra*, and

Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding XMES, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, modifications of gene expression can be achieved by designing complementary sequences or antisense molecules (DNA, RNA, PNA, or modified oligonucleotides) to the coding or regulatory regions of the gene encoding XMES. Such technology is well known in the art, and antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding XMES. (See, e.g., Agrawal, S., ed. (1996) Antisense Therapeutics, Humana Press Inc., Totawa NJ.)

In therapeutic use, any gene delivery system suitable for introduction of the antisense sequences into appropriate target cells can be used. Antisense sequences can be delivered intracellularly in the form of an expression plasmid which, upon transcription, produces a sequence complementary to at least a portion of the cellular sequence encoding the target protein. (See, e.g., Slater, J.E. et al. (1998) *J. Allergy Clin. Immunol.* 102(3):469-475; and Scanlon, K.J. et al. (1995) 9(13):1288-1296.) Antisense sequences can also be introduced intracellularly through the use of viral vectors, such as retrovirus and adeno-associated virus vectors. (See, e.g., Miller, A.D. (1990) *Blood* 76:271; Ausubel, supra; Uckert, W. and W. Walther (1994) *Pharmacol. Ther.* 63(3):323-347.) Other gene delivery mechanisms include liposome-derived systems, artificial viral envelopes, and other systems known in the art. (See, e.g., Rossi, J.J. (1995) *Br. Med. Bull.* 51(1):217-225; Boado, R.J. et al. (1998) *J. Pharm. Sci.* 87(11):1308-1315; and Morris, M.C. et al. (1997) *Nucleic Acids Res.* 25(14):2730-2736.)

In another embodiment of the invention, polynucleotides encoding XMES may be used for somatic or germline gene therapy. Gene therapy may be performed to (i) correct a genetic deficiency (e.g., in the cases of severe combined immunodeficiency (SCID)-X1 disease characterized by X-linked inheritance (Cavazzana-Calvo, M. et al. (2000) *Science* 288:669-672), severe combined immunodeficiency syndrome associated with an inherited adenosine deaminase (ADA) deficiency (Blaese, R.M. et al. (1995) *Science* 270:475-480; Bordignon, C. et al. (1995) *Science* 270:470-475), cystic fibrosis (Zabner, J. et al. (1993) *Cell* 75:207-216; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:643-666; Crystal, R.G. et al. (1995) *Hum. Gene Therapy* 6:667-703), thalassemias, familial hypercholesterolemia, and hemophilia resulting from Factor VIII or Factor IX deficiencies (Crystal, R.G. (1995) *Science* 270:404-410; Verma, I.M. and N. Somia (1997) *Nature* 389:239-242)), (ii) express a conditionally lethal gene product (e.g., in the case of cancers which result from unregulated cell proliferation), or (iii) express a protein which affords protection against intracellular parasites (e.g., against human retroviruses, such as human immunodeficiency virus (HIV) (Baltimore, D. (1988) *Nature* 335:395-396; Poeschla, E. et al. (1996) *Proc. Natl. Acad. Sci. USA.* 93:11395-11399),

hepatitis B or C virus (HBV, HCV); fungal parasites, such as Candida albicans and Paracoccidioides brasiliensis; and protozoan parasites such as Plasmodium falciparum and Trypanosoma cruzi). In the case where a genetic deficiency in XMES expression or regulation causes disease, the expression of XMES from an appropriate population of transduced cells may alleviate the clinical manifestations caused by the genetic deficiency.

In a further embodiment of the invention, diseases or disorders caused by deficiencies in XMES are treated by constructing mammalian expression vectors encoding XMES and introducing these vectors by mechanical means into XMES-deficient cells. Mechanical transfer technologies for use with cells in vivo or ex vitro include (i) direct DNA microinjection into individual cells, (ii) ballistic gold particle delivery, (iii) liposome-mediated transfection, (iv) receptor-mediated gene transfer, and (v) the use of DNA transposons (Morgan, R.A. and W.F. Anderson (1993) *Annu. Rev. Biochem.* 62:191-217; Ivics, Z. (1997) *Cell* 91:501-510; Boulay, J-L. and H. Récipon (1998) *Curr. Opin. Biotechnol.* 9:445-450).

Expression vectors that may be effective for the expression of XMES include, but are not limited to, the PCDNA 3.1, EPITAG, PRCCMV2, PREP, PVAX vectors (Invitrogen, Carlsbad CA), PCMV-SCRIPT, PCMV-TAG, PEGSH/PERV (Stratagene, La Jolla CA), and PTET-OFF, PTET-ON, PTRE2, PTRE2-LUC, PTK-HYG (Clontech, Palo Alto CA). XMES may be expressed using (i) a constitutively active promoter, (e.g., from cytomegalovirus (CMV), Rous sarcoma virus (RSV), SV40 virus, thymidine kinase (TK), or β -actin genes), (ii) an inducible promoter (e.g., the tetracycline-regulated promoter (Gossen, M. and H. Bujard (1992) *Proc. Natl. Acad. Sci. USA* 89:5547-5551; Gossen, M. et al. (1995) *Science* 268:1766-1769; Rossi, F.M.V. and H.M. Blau (1998) *Curr. Opin. Biotechnol.* 9:451-456), commercially available in the T-REX plasmid (Invitrogen)); the ecdysone-inducible promoter (available in the plasmids PVGRXR and PIND; Invitrogen); the FK506/rapamycin inducible promoter; or the RU486/mifepristone inducible promoter (Rossi, F.M.V. and Blau, H.M. *supra*)), or (iii) a tissue-specific promoter or the native promoter of the endogenous gene encoding XMES from a normal individual.

Commercially available liposome transformation kits (e.g., the PERFECT LIPID TRANSFECTION KIT, available from Invitrogen) allow one with ordinary skill in the art to deliver polynucleotides to target cells in culture and require minimal effort to optimize experimental parameters. In the alternative, transformation is performed using the calcium phosphate method (Graham, F.L. and A.J. Eb (1973) *Virology* 52:456-467), or by electroporation (Neumann, E. et al. (1982) *EMBO J.* 1:841-845). The introduction of DNA to primary cells requires modification of these standardized mammalian transfection protocols.

In another embodiment of the invention, diseases or disorders caused by genetic defects with

- respect to XMES expression are treated by constructing a retrovirus vector consisting of (i) the polynucleotide encoding XMES under the control of an independent promoter or the retrovirus long terminal repeat (LTR) promoter, (ii) appropriate RNA packaging signals, and (iii) a Rev-responsive element (RRE) along with additional retrovirus *cis*-acting RNA sequences and coding sequences
- 5 required for efficient vector propagation. Retrovirus vectors (e.g., PFB and PFBNEO) are commercially available (Stratagene) and are based on published data (Riviere, I. et al. (1995) Proc. Natl. Acad. Sci. USA 92:6733-6737), incorporated by reference herein. The vector is propagated in an appropriate vector producing cell line (VPCL) that expresses an envelope gene with a tropism for receptors on the target cells or a promiscuous envelope protein such as VSVg (Armentano, D. et al.
- 10 (1987) J. Virol. 61:1647-1650; Bender, M.A. et al. (1987) J. Virol. 61:1639-1646; Adam, M.A. and A.D. Miller (1988) J. Virol. 62:3802-3806; Dull, T. et al. (1998) J. Virol. 72:8463-8471; Zufferey, R. et al. (1998) J. Virol. 72:9873-9880). U.S. Patent Number 5,910,434 to Rigg ("Method for obtaining retrovirus packaging cell lines producing high transducing efficiency retroviral supernatant") discloses a method for obtaining retrovirus packaging cell lines and is hereby incorporated by reference.
- 15 Propagation of retrovirus vectors, transduction of a population of cells (e.g., CD4⁺ T-cells), and the return of transduced cells to a patient are procedures well known to persons skilled in the art of gene therapy and have been well documented (Ranga, U. et al. (1997) J. Virol. 71:7020-7029; Bauer, G. et al. (1997) Blood 89:2259-2267; Bonyhadi, M.L. (1997) J. Virol. 71:4707-4716; Ranga, U. et al. (1998) Proc. Natl. Acad. Sci. USA 95:1201-1206; Su, L. (1997) Blood 89:2283-2290).
- 20 In the alternative, an adenovirus-based gene therapy delivery system is used to deliver polynucleotides encoding XMES to cells which have one or more genetic abnormalities with respect to the expression of XMES. The construction and packaging of adenovirus-based vectors are well known to those with ordinary skill in the art. Replication defective adenovirus vectors have proven to be versatile for importing genes encoding immunoregulatory proteins into intact islets in the pancreas
- 25 (Csete, M.E. et al. (1995) Transplantation 27:263-268). Potentially useful adenoviral vectors are described in U.S. Patent Number 5,707,618 to Armentano ("Adenovirus vectors for gene therapy"), hereby incorporated by reference. For adenoviral vectors, see also Antinozzi, P.A. et al. (1999) Annu. Rev. Nutr. 19:511-544 and Verma, I.M. and N. Somia (1997) Nature 389:239-242, both incorporated by reference herein.
- 30 In another alternative, a herpes-based, gene therapy delivery system is used to deliver polynucleotides encoding XMES to target cells which have one or more genetic abnormalities with respect to the expression of XMES. The use of herpes simplex virus (HSV)-based vectors may be especially valuable for introducing XMES to cells of the central nervous system, for which HSV has a tropism. The construction and packaging of herpes-based vectors are well known to those with

ordinary skill in the art. A replication-competent herpes simplex virus (HSV) type 1-based vector has been used to deliver a reporter gene to the eyes of primates (Liu, X. et al. (1999) Exp. Eye Res. 169:385-395). The construction of a HSV-1 virus vector has also been disclosed in detail in U.S. Patent Number 5,804,413 to DeLuca ("Herpes simplex virus strains for gene transfer"), which is hereby incorporated by reference. U.S. Patent Number 5,804,413 teaches the use of recombinant HSV d92 which consists of a genome containing at least one exogenous gene to be transferred to a cell under the control of the appropriate promoter for purposes including human gene therapy. Also taught by this patent are the construction and use of recombinant HSV strains deleted for ICP4, ICP27 and ICP22. For HSV vectors, see also Goins, W.F. et al. (1999) J. Virol. 73:519-532 and Xu, H. et al. (1994) Dev. Biol. 163:152-161, hereby incorporated by reference. The manipulation of cloned herpesvirus sequences, the generation of recombinant virus following the transfection of multiple plasmids containing different segments of the large herpesvirus genomes, the growth and propagation of herpesvirus, and the infection of cells with herpesvirus are techniques well known to those of ordinary skill in the art.

In another alternative, an alphavirus (positive, single-stranded RNA virus) vector is used to deliver polynucleotides encoding XMES to target cells. The biology of the prototypic alphavirus, Semliki Forest Virus (SFV), has been studied extensively and gene transfer vectors have been based on the SFV genome (Garoff, H. and K.-J. Li (1998) Curr. Opin. Biotechnol. 9:464-469). During alphavirus RNA replication, a subgenomic RNA is generated that normally encodes the viral capsid proteins. This subgenomic RNA replicates to higher levels than the full length genomic RNA, resulting in the overproduction of capsid proteins relative to the viral proteins with enzymatic activity (e.g., protease and polymerase). Similarly, inserting the coding sequence for XMES into the alphavirus genome in place of the capsid-coding region results in the production of a large number of XMES-coding RNAs and the synthesis of high levels of XMES in vector transduced cells. While alphavirus infection is typically associated with cell lysis within a few days, the ability to establish a persistent infection in hamster normal kidney cells (BHK-21) with a variant of Sindbis virus (SIN) indicates that the lytic replication of alphaviruses can be altered to suit the needs of the gene therapy application (Dryga, S.A. et al. (1997) Virology 228:74-83). The wide host range of alphaviruses will allow the introduction of XMES into a variety of cell types. The specific transduction of a subset of cells in a population may require the sorting of cells prior to transduction. The methods of manipulating infectious cDNA clones of alphaviruses, performing alphavirus cDNA and RNA transfections, and performing alphavirus infections, are well known to those with ordinary skill in the art.

Oligonucleotides derived from the transcription initiation site, e.g., between about positions -10 and +10 from the start site, may also be employed to inhibit gene expression. Similarly, inhibition can

be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber, B.E. and B.I. Carr, 5 Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme 10 molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding XMES.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA, 15 GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides, corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

20 Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding XMES. Such DNA sequences may be incorporated into a wide variety of vectors with 25 suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends 30 of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

An additional embodiment of the invention encompasses a method for screening for a compound which is effective in altering expression of a polynucleotide encoding XMES.

Compounds which may be effective in altering expression of a specific polynucleotide may include, but are not limited to, oligonucleotides, antisense oligonucleotides, triple helix-forming

5 oligonucleotides, transcription factors and other polypeptide transcriptional regulators, and non-macromolecular chemical entities which are capable of interacting with specific polynucleotide sequences. Effective compounds may alter polynucleotide expression by acting as either inhibitors or promoters of polynucleotide expression. Thus, in the treatment of disorders associated with increased XMES expression or activity, a compound which specifically inhibits expression of the
10 polynucleotide encoding XMES may be therapeutically useful, and in the treatment of disorders associated with decreased XMES expression or activity, a compound which specifically promotes expression of the polynucleotide encoding XMES may be therapeutically useful.

At least one, and up to a plurality, of test compounds may be screened for effectiveness in altering expression of a specific polynucleotide. A test compound may be obtained by any method
15 commonly known in the art, including chemical modification of a compound known to be effective in altering polynucleotide expression; selection from an existing, commercially-available or proprietary library of naturally-occurring or non-natural chemical compounds; rational design of a compound based on chemical and/or structural properties of the target polynucleotide; and selection from a library of chemical compounds created combinatorially or randomly. A sample comprising a
20 polynucleotide encoding XMES is exposed to at least one test compound thus obtained. The sample may comprise, for example, an intact or permeabilized cell, or an *in vitro* cell-free or reconstituted biochemical system. Alterations in the expression of a polynucleotide encoding XMES are assayed by any method commonly known in the art. Typically, the expression of a specific nucleotide is detected by hybridization with a probe having a nucleotide sequence complementary to the sequence
25 of the polynucleotide encoding XMES. The amount of hybridization may be quantified, thus forming the basis for a comparison of the expression of the polynucleotide both with and without exposure to one or more test compounds. Detection of a change in the expression of a polynucleotide exposed to a test compound indicates that the test compound is effective in altering the expression of the polynucleotide. A screen for a compound effective in altering expression of a specific
30 polynucleotide can be carried out, for example, using a *Schizosaccharomyces pombe* gene expression system (Atkins, D. et al. (1999) U.S. Patent No. 5,932,435; Arndt, G.M. et al. (2000) Nucleic Acids Res. 28:E15) or a human cell line such as HeLa cell (Clarke, M.L. et al. (2000) Biochem. Biophys. Res. Commun. 268:8-13). A particular embodiment of the present invention involves screening a combinatorial library of oligonucleotides (such as deoxyribonucleotides, ribonucleotides, peptide
35 nucleic acids, and modified oligonucleotides) for antisense activity against a specific polynucleotide

sequence (Bruice, T.W. et al. (1997) U.S. Patent No. 5,686,242; Bruice, T.W. et al. (2000) U.S. Patent No. 6,022,691).

Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat. Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a composition which generally comprises an active ingredient formulated with a pharmaceutically acceptable excipient. Excipients may include, for example, sugars, starches, celluloses, gums, and proteins. Various formulations are commonly known and are thoroughly discussed in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing, Easton PA). Such compositions may consist of XMES, antibodies to XMES, and mimetics, agonists, antagonists, or inhibitors of XMES.

The compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, pulmonary, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

Compositions for pulmonary administration may be prepared in liquid or dry powder form. These compositions are generally aerosolized immediately prior to inhalation by the patient. In the case of small molecules (e.g. traditional low molecular weight organic drugs), aerosol delivery of fast-acting formulations is well-known in the art. In the case of macromolecules (e.g. larger peptides and proteins), recent developments in the field of pulmonary delivery via the alveolar region of the lung have enabled the practical delivery of drugs such as insulin to blood circulation (see, e.g., Patton, J.S. et al., U.S. Patent No. 5,997,848). Pulmonary delivery has the advantage of administration without needle injection, and obviates the need for potentially toxic penetration enhancers.

Compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

Specialized forms of compositions may be prepared for direct intracellular delivery of macromolecules comprising XMES or fragments thereof. For example, liposome preparations

containing a cell-impermeable macromolecule may promote cell fusion and intracellular delivery of the macromolecule. Alternatively, XMES or a fragment thereof may be joined to a short cationic N-terminal portion from the HIV Tat-1 protein. Fusion proteins thus generated have been found to transduce into the cells of all tissues, including the brain, in a mouse model system (Schwarze, S.R. et al. (1999) Science 285:1569-1572).

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, monkeys, or pigs. An animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example XMES or fragments thereof, antibodies of XMES, and agonists, antagonists or inhibitors of XMES, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such as by calculating the ED₅₀ (the dose therapeutically effective in 50% of the population) or LD₅₀ (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD₅₀/ED₅₀ ratio. Compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED₅₀ with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 µg to 100,000 µg, up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells,

conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind XMES may be used for the diagnosis of disorders characterized by expression of XMES, or in assays to monitor patients being
5 treated with XMES or agonists, antagonists, or inhibitors of XMES. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for XMES include methods which utilize the antibody and a label to detect XMES in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter
10 molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring XMES, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of XMES expression. Normal or standard values for XMES expression are established by combining body fluids or cell extracts taken from normal mammalian subjects, for example, human subjects, with antibodies to XMES under
15 conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of XMES expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values. Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding XMES may be used for
20 diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of XMES may be correlated with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of XMES, and to monitor regulation of XMES levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding XMES or closely related molecules may be used to identify nucleic acid sequences which encode XMES. The specificity of the probe, whether it is made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the
30 probe identifies only naturally occurring sequences encoding XMES, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50% sequence identity to any of the XMES encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:10-18 or from

genomic sequences including promoters, enhancers, and introns of the XMES gene.

Means for producing specific hybridization probes for DNAs encoding XMES include the cloning of polynucleotide sequences encoding XMES or XMES derivatives into vectors for the production of mRNA probes. Such vectors are known in the art, are commercially available, and may
5 be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

Polynucleotide sequences encoding XMES may be used for the diagnosis of disorders
10 associated with expression of XMES. Examples of such disorders include, but are not limited to, a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other
15 demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal
20 syndrome, mental retardation and other developmental disorders of the central nervous system including Down syndrome, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety,
25 and schizophrenic disorders, seasonal affective disorder (SAD), akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, Tourette's disorder, progressive supranuclear palsy, corticobasal degeneration, and familial frontotemporal dementia; an autoimmune/inflammatory disorder such as acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis,
30 anemia, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy (APECED), bronchitis, cholecystitis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, episodic lymphopenia with lymphocytotoxins, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis,

hypereosinophilia, irritable bowel syndrome, multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, osteoarthritis, osteoporosis, pancreatitis, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, thrombocytopenic purpura, ulcerative colitis, uveitis, Werner

5 syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, viral, bacterial, fungal, parasitic, protozoal, and helminthic infections, and trauma; a developmental disorder such as renal tubular acidosis, anemia, Cushing's syndrome, achondroplastic dwarfism, Duchenne and Becker muscular dystrophy, epilepsy, gonadal dysgenesis, WAGR syndrome (Wilms' tumor, aniridia, genitourinary abnormalities, and mental retardation), Smith-Magenis syndrome, myelodysplastic

10 syndrome, hereditary mucoepithelial dysplasia, hereditary keratodermas, hereditary neuropathies such as Charcot-Marie-Tooth disease and neurofibromatosis, hypothyroidism, hydrocephalus, seizure disorders such as Sydenham's chorea and cerebral palsy, spina bifida, anencephaly, craniorachischisis, congenital glaucoma, cataract, and sensorineural hearing loss; an endocrine disorder such as a disorder of the hypothalamus and/or pituitary resulting from lesions such as a primary brain tumor, adenoma,

15 infarction associated with pregnancy, hypophysectomy, aneurysm, vascular malformation, thrombosis, infection, immunological disorder, and complication due to head trauma; a disorder associated with hypopituitarism including hypogonadism, Sheehan syndrome, diabetes insipidus, Kallman's disease, Hand-Schuller-Christian disease, Letterer-Siwe disease, sarcoidosis, empty sella syndrome, and dwarfism; a disorder associated with hyperpituitarism including acromegaly, gigantism, and syndrome of

20 inappropriate antidiuretic hormone (ADH) secretion (SIADH) often caused by benign adenoma; a disorder associated with hypothyroidism including goiter, myxedema, acute thyroiditis associated with bacterial infection, subacute thyroiditis associated with viral infection, autoimmune thyroiditis (Hashimoto's disease), and cretinism; a disorder associated with hyperthyroidism including thyrotoxicosis and its various forms, Grave's disease, pretibial myxedema, toxic multinodular goiter,

25 thyroid carcinoma, and Plummer's disease; a disorder associated with hyperparathyroidism including Conn disease (chronic hypercalcemia); a pancreatic disorder such as Type I or Type II diabetes mellitus and associated complications; a disorder associated with the adrenals such as hyperplasia, carcinoma, or adenoma of the adrenal cortex, hypertension associated with alkalosis, amyloidosis, hypokalemia, Cushing's disease, Liddle's syndrome, and Arnold-Healy-Gordon syndrome, pheochromocytoma

30 tumors, and Addison's disease; a disorder associated with gonadal steroid hormones such as: in women, abnormal prolactin production, infertility, endometriosis, perturbation of the menstrual cycle, polycystic ovarian disease, hyperprolactinemia, isolated gonadotropin deficiency, amenorrhea, galactorrhea, hermaphroditism, hirsutism and virilization, breast cancer, and, in post-menopausal women, osteoporosis; and, in men, Leydig cell deficiency, male climacteric phase, and germinal cell

aplasia, a hypergonadal disorder associated with Leydig cell tumors, androgen resistance associated with absence of androgen receptors, 5 α -reductase deficiency, 21-hydroxylase deficiency, and gynecomastia, and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, 5 paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

10 The polynucleotide sequences encoding XMES may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered XMES expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding XMES may be useful in assays that 15 detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding XMES may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control 20 sample then the presence of altered levels of nucleotide sequences encoding XMES in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of XMES, 25 a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding XMES, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used.

30 Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the

patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

Additional diagnostic uses for oligonucleotides designed from the sequences encoding XMES may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding XMES, or a fragment of a polynucleotide complementary to the polynucleotide encoding XMES, and will be employed under optimized conditions for identification of a specific gene or condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

In a particular aspect, oligonucleotide primers derived from the polynucleotide sequences encoding XMES may be used to detect single nucleotide polymorphisms (SNPs). SNPs are substitutions, insertions and deletions that are a frequent cause of inherited or acquired genetic disease in humans. Methods of SNP detection include, but are not limited to, single-stranded conformation polymorphism (SSCP) and fluorescent SSCP (fSSCP) methods. In SSCP, oligonucleotide primers derived from the polynucleotide sequences encoding XMES are used to amplify DNA using the polymerase chain reaction (PCR). The DNA may be derived, for example, from diseased or normal tissue, biopsy samples, bodily fluids, and the like. SNPs in the DNA cause differences in the secondary and tertiary structures of PCR products in single-stranded form, and these differences are detectable using gel electrophoresis in non-denaturing gels. In fSSCP, the oligonucleotide primers are fluorescently labeled, which allows detection of the amplimers in high-throughput equipment such as DNA sequencing machines. Additionally, sequence database analysis methods, termed *in silico* SNP (isSNP), are capable of identifying polymorphisms by comparing the sequence of individual overlapping DNA fragments which assemble into a common consensus sequence. These computer-based methods filter out sequence variations due to laboratory preparation of DNA and sequencing errors using statistical models and automated analyses of DNA sequence chromatograms. In the alternative, SNPs may be detected and characterized by mass spectrometry using, for example, the high throughput MASSARRAY system (Sequenom, Inc., San Diego CA).

Methods which may also be used to quantify the expression of XMES include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C. et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be
5 accelerated by running the assay in a high-throughput format where the oligomer or polynucleotide of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the polynucleotide sequences described herein may be used as elements on a microarray. The microarray
10 can be used in transcript imaging techniques which monitor the relative expression levels of large numbers of genes simultaneously as described below. The microarray may also be used to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, to monitor progression/regression of disease as a function of gene expression, and to develop and monitor the
15 activities of therapeutic agents in the treatment of disease. In particular, this information may be used to develop a pharmacogenomic profile of a patient in order to select the most appropriate and effective treatment regimen for that patient. For example, therapeutic agents which are highly effective and display the fewest side effects may be selected for a patient based on his/her pharmacogenomic profile.

In another embodiment, XMES, fragments of XMES, or antibodies specific for XMES may be
20 used as elements on a microarray. The microarray may be used to monitor or measure protein-protein interactions, drug-target interactions, and gene expression profiles, as described above.

A particular embodiment relates to the use of the polynucleotides of the present invention to generate a transcript image of a tissue or cell type. A transcript image represents the global pattern of gene expression by a particular tissue or cell type. Global gene expression patterns are analyzed by
25 quantifying the number of expressed genes and their relative abundance under given conditions and at a given time. (See Seilhamer et al., "Comparative Gene Transcript Analysis," U.S. Patent Number 5,840,484, expressly incorporated by reference herein.) Thus a transcript image may be generated by hybridizing the polynucleotides of the present invention or their complements to the totality of transcripts or reverse transcripts of a particular tissue or cell type. In one embodiment, the
30 hybridization takes place in high-throughput format, wherein the polynucleotides of the present invention or their complements comprise a subset of a plurality of elements on a microarray. The resultant transcript image would provide a profile of gene activity.

Transcript images may be generated using transcripts isolated from tissues, cell lines, biopsies, or other biological samples. The transcript image may thus reflect gene expression in vivo, as in the

case of a tissue or biopsy sample, or in vitro, as in the case of a cell line.

Transcript images which profile the expression of the polynucleotides of the present invention may also be used in conjunction with in vitro model systems and preclinical evaluation of pharmaceuticals, as well as toxicological testing of industrial and naturally-occurring environmental compounds. All compounds induce characteristic gene expression patterns, frequently termed molecular fingerprints or toxicant signatures, which are indicative of mechanisms of action and toxicity (Nuwaysir, E.F. et al. (1999) Mol. Carcinog. 24:153-159; Steiner, S. and N.L. Anderson (2000) Toxicol. Lett. 112-113:467-471, expressly incorporated by reference herein). If a test compound has a signature similar to that of a compound with known toxicity, it is likely to share those toxic properties. These fingerprints or signatures are most useful and refined when they contain expression information from a large number of genes and gene families. Ideally, a genome-wide measurement of expression provides the highest quality signature. Even genes whose expression is not altered by any tested compounds are important as well, as the levels of expression of these genes are used to normalize the rest of the expression data. The normalization procedure is useful for comparison of expression data after treatment with different compounds. While the assignment of gene function to elements of a toxicant signature aids in interpretation of toxicity mechanisms, knowledge of gene function is not necessary for the statistical matching of signatures which leads to prediction of toxicity. (See, for example, Press Release 00-02 from the National Institute of Environmental Health Sciences, released February 29, 2000, available at <http://www.niehs.nih.gov/oc/news/toxchip.htm>.) Therefore, it is important and desirable in toxicological screening using toxicant signatures to include all expressed gene sequences.

In one embodiment, the toxicity of a test compound is assessed by treating a biological sample containing nucleic acids with the test compound. Nucleic acids that are expressed in the treated biological sample are hybridized with one or more probes specific to the polynucleotides of the present invention, so that transcript levels corresponding to the polynucleotides of the present invention may be quantified. The transcript levels in the treated biological sample are compared with levels in an untreated biological sample. Differences in the transcript levels between the two samples are indicative of a toxic response caused by the test compound in the treated sample.

Another particular embodiment relates to the use of the polypeptide sequences of the present invention to analyze the proteome of a tissue or cell type. The term proteome refers to the global pattern of protein expression in a particular tissue or cell type. Each protein component of a proteome can be subjected individually to further analysis. Proteome expression patterns, or profiles, are analyzed by quantifying the number of expressed proteins and their relative abundance under given conditions and at a given time. A profile of a cell's proteome may thus be generated by separating and

analyzing the polypeptides of a particular tissue or cell type. In one embodiment, the separation is achieved using two-dimensional gel electrophoresis, in which proteins from a sample are separated by isoelectric focusing in the first dimension, and then according to molecular weight by sodium dodecyl sulfate slab gel electrophoresis in the second dimension (Steiner and Anderson, supra). The proteins are
5 visualized in the gel as discrete and uniquely positioned spots, typically by staining the gel with an agent such as Coomassie Blue or silver or fluorescent stains. The optical density of each protein spot is generally proportional to the level of the protein in the sample. The optical densities of equivalently positioned protein spots from different samples, for example, from biological samples either treated or untreated with a test compound or therapeutic agent, are compared to identify any changes in protein
10 spot density related to the treatment. The proteins in the spots are partially sequenced using, for example, standard methods employing chemical or enzymatic cleavage followed by mass spectrometry. The identity of the protein in a spot may be determined by comparing its partial sequence, preferably of at least 5 contiguous amino acid residues, to the polypeptide sequences of the present invention. In some cases, further sequence data may be obtained for definitive protein identification.

15 A proteomic profile may also be generated using antibodies specific for XMES to quantify the levels of XMES expression. In one embodiment, the antibodies are used as elements on a microarray, and protein expression levels are quantified by exposing the microarray to the sample and detecting the levels of protein bound to each array element (Lueking, A. et al. (1999) *Anal. Biochem.* 270:103-111; Mendoz, L.G. et al. (1999) *Biotechniques* 27:778-788). Detection may be performed by a variety of
20 methods known in the art, for example, by reacting the proteins in the sample with a thiol- or amino-reactive fluorescent compound and detecting the amount of fluorescence bound at each array element.

Toxicant signatures at the proteome level are also useful for toxicological screening, and should be analyzed in parallel with toxicant signatures at the transcript level. There is a poor correlation between transcript and protein abundances for some proteins in some tissues (Anderson, N.L. and J.
25 Seilhamer (1997) *Electrophoresis* 18:533-537), so proteome toxicant signatures may be useful in the analysis of compounds which do not significantly affect the transcript image, but which alter the proteomic profile. In addition, the analysis of transcripts in body fluids is difficult, due to rapid degradation of mRNA, so proteomic profiling may be more reliable and informative in such cases.

In another embodiment, the toxicity of a test compound is assessed by treating a biological
30 sample containing proteins with the test compound. Proteins that are expressed in the treated biological sample are separated so that the amount of each protein can be quantified. The amount of each protein is compared to the amount of the corresponding protein in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample. Individual proteins are identified by sequencing the amino acid residues of the

individual proteins and comparing these partial sequences to the polypeptides of the present invention.

In another embodiment, the toxicity of a test compound is assessed by treating a biological sample containing proteins with the test compound. Proteins from the biological sample are incubated with antibodies specific to the polypeptides of the present invention. The amount of protein recognized by the antibodies is quantified. The amount of protein in the treated biological sample is compared with the amount in an untreated biological sample. A difference in the amount of protein between the two samples is indicative of a toxic response to the test compound in the treated sample.

Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.) Various types of microarrays are well known and thoroughly described in DNA Microarrays: A Practical Approach, M. Schena, ed. (1999) Oxford University Press, London, hereby expressly incorporated by reference.

In another embodiment of the invention, nucleic acid sequences encoding XMES may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. Either coding or noncoding sequences may be used, and in some instances, noncoding sequences may be preferable over coding sequences. For example, conservation of a coding sequence among members of a multi-gene family may potentially cause undesired cross hybridization during chromosomal mapping. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price, C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.) Once mapped, the nucleic acid sequences of the invention may be used to develop genetic linkage maps, for example, which correlate the inheritance of a disease state with the inheritance of a particular chromosome region or restriction fragment length polymorphism (RFLP). (See, for example, Lander, E.S. and D. Botstein (1986) Proc. Natl. Acad. Sci. USA 83:7353-7357.)

Fluorescent *in situ* hybridization (FISH) may be correlated with other physical and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, *supra*, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding XMES on a physical map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder and thus may further positional cloning efforts.

In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the exact chromosomal locus is not known. This information is
5 valuable to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the gene or genes responsible for a disease or syndrome have been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the instant invention may
10 also be used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, XMES, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a
15 solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between XMES and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are
20 synthesized on a solid substrate. The test compounds are reacted with XMES, or fragments thereof, and washed. Bound XMES is then detected by methods well known in the art. Purified XMES can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

25 In another embodiment, one may use competitive drug screening assays in which neutralizing antibodies capable of binding XMES specifically compete with a test compound for binding XMES. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with XMES.

In additional embodiments, the nucleotide sequences which encode XMES may be used in any
30 molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are,

therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications and publications, mentioned above and below, including U.S. Ser. No. 60/210,233, U.S. Ser. No. 60/213,465, and U.S. Ser. No. 60/249,019 are
5 expressly incorporated by reference herein.

EXAMPLES

I. Construction of cDNA Libraries

Incyte cDNAs were derived from cDNA libraries described in the LIFESEQ GOLD database
10 (Incyte Genomics, Palo Alto CA) and shown in Table 4, column 5. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and
15 ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A)+ RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was
20 isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the
25 recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column
30 chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUESCRIPT plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), PCDNA2.1 plasmid (Invitrogen, Carlsbad CA), PBK-CMV plasmid (Stratagene), or pINCY (Incyte Genomics, Palo Alto CA), or derivatives thereof. Recombinant plasmids were transformed into competent E. coli cells

including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids obtained as described in Example I were recovered from host cells by *in vivo* excision
5 using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of
10 distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using
15 PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

Incyte cDNA recovered in plasmids as described in Example II were sequenced as follows. Sequencing reactions were processed using standard methods or high-throughput instrumentation
20 such as the ABI CATALYST 800 (Applied Biosystems) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Applied Biosystems).
25 Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the ABI PRISM 373 or 377 sequencing system (Applied Biosystems) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997,
30 *supra*, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example VIII.

The polynucleotide sequences derived from Incyte cDNAs were validated by removing vector, linker, and poly(A) sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The Incyte cDNA

sequences or translations thereof were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and hidden Markov model (HMM)-based protein family databases such as PFAM. (HMM is a probabilistic approach which analyzes consensus primary structures of gene families.

- 5 See, for example, Eddy, S.R. (1996) *Curr. Opin. Struct. Biol.* 6:361-365.) The queries were performed using programs based on BLAST, FASTA, BLIMPS, and HMMER. The Incyte cDNA sequences were assembled to produce full length polynucleotide sequences. Alternatively, GenBank cDNAs, GenBank ESTs, stitched sequences, stretched sequences, or Genscan-predicted coding sequences (see Examples IV and V) were used to extend Incyte cDNA assemblages to full length.
- 10 Assembly was performed using programs based on Phred, Phrap, and Consed, and cDNA assemblages were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length polypeptide sequences. Alternatively, a polypeptide of the invention may begin at any of the methionine residues of the full length translated polypeptide. Full length polypeptide sequences were subsequently
- 15 analyzed by querying against databases such as the GenBank protein databases (genpept), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and hidden Markov model (HMM)-based protein family databases such as PFAM. Full length polynucleotide sequences are also analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments are
- 20 generated using default parameters specified by the CLUSTAL algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

- Table 7 summarizes the tools, programs, and algorithms used for the analysis and assembly of Incyte cDNA and full length sequences and provides applicable descriptions, references, and threshold
- 25 parameters. The first column of Table 7 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score or the lower the probability value, the greater the identity
- 30 between two sequences).

The programs described above for the assembly and analysis of full length polynucleotide and polypeptide sequences were also used to identify polynucleotide sequence fragments from SEQ ID NO:10-18. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies are described in Table 4, column 4.

IV. Identification and Editing of Coding Sequences from Genomic DNA

Putative extracellular messengers were initially identified by running the Genscan gene identification program against public genomic sequence databases (e.g., gbpri and gbhtg). Genscan is a general-purpose gene identification program which analyzes genomic DNA sequences from a variety of organisms (See Burge, C. and S. Karlin (1997) J. Mol. Biol. 268:78-94, and Burge, C. and S. Karlin (1998) Curr. Opin. Struct. Biol. 8:346-354). The program concatenates predicted exons to form an assembled cDNA sequence extending from a methionine to a stop codon. The output of Genscan is a FASTA database of polynucleotide and polypeptide sequences. The maximum range of sequence for Genscan to analyze at once was set to 30 kb. To determine which of these Genscan predicted cDNA sequences encode extracellular messengers, the encoded polypeptides were analyzed by querying against PFAM models for extracellular messengers. Potential extracellular messengers were also identified by homology to Incyte cDNA sequences that had been annotated as extracellular messengers. These selected Genscan-predicted sequences were then compared by BLAST analysis to the genpept and gbpri public databases. Where necessary, the Genscan-predicted sequences were then edited by comparison to the top BLAST hit from genpept to correct errors in the sequence predicted by Genscan, such as extra or omitted exons. BLAST analysis was also used to find any Incyte cDNA or public cDNA coverage of the Genscan-predicted sequences, thus providing evidence for transcription. When Incyte cDNA coverage was available, this information was used to correct or confirm the Genscan predicted sequence. Full length polynucleotide sequences were obtained by assembling Genscan-predicted coding sequences with Incyte cDNA sequences and/or public cDNA sequences using the assembly process described in Example III. Alternatively, full length polynucleotide sequences were derived entirely from edited or unedited Genscan-predicted coding sequences.

V. Assembly of Genomic Sequence Data with cDNA Sequence Data

"Stitched" Sequences

Partial cDNA sequences were extended with exons predicted by the Genscan gene identification program described in Example IV. Partial cDNAs assembled as described in Example III were mapped to genomic DNA and parsed into clusters containing related cDNAs and Genscan exon predictions from one or more genomic sequences. Each cluster was analyzed using an algorithm based on graph theory and dynamic programming to integrate cDNA and genomic information, generating possible splice variants that were subsequently confirmed, edited, or extended to create a full length sequence. Sequence intervals in which the entire length of the interval was present on more than one sequence in the cluster were identified, and intervals thus identified were considered to be equivalent by transitivity. For example, if an interval was present on a cDNA and two genomic sequences, then all three intervals were considered to be equivalent. This process allows unrelated but consecutive genomic sequences to

be brought together, bridged by cDNA sequence. Intervals thus identified were then "stitched" together by the stitching algorithm in the order that they appear along their parent sequences to generate the longest possible sequence, as well as sequence variants. Linkages between intervals which proceed along one type of parent sequence (cDNA to cDNA or genomic sequence to genomic sequence) were given preference over linkages which change parent type (cDNA to genomic sequence). The resultant stitched sequences were translated and compared by BLAST analysis to the genpept and gbpri public databases. Incorrect exons predicted by Genscan were corrected by comparison to the top BLAST hit from genpept. Sequences were further extended with additional cDNA sequences, or by inspection of genomic DNA, when necessary.

10 **"Stretched" Sequences**

Partial DNA sequences were extended to full length with an algorithm based on BLAST analysis. First, partial cDNAs assembled as described in Example III were queried against public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases using the BLAST program. The nearest GenBank protein homolog was then compared by BLAST analysis to either Incyte cDNA sequences or GenScan exon predicted sequences described in Example IV. A chimeric protein was generated by using the resultant high-scoring segment pairs (HSPs) to map the translated sequences onto the GenBank protein homolog. Insertions or deletions may occur in the chimeric protein with respect to the original GenBank protein homolog. The GenBank protein homolog, the chimeric protein, or both were used as probes to search for homologous genomic sequences from the public human genome databases. Partial DNA sequences were therefore "stretched" or extended by the addition of homologous genomic sequences. The resultant stretched sequences were examined to determine whether it contained a complete gene.

VI. Chromosomal Mapping of XMES Encoding Polynucleotides

The sequences which were used to assemble SEQ ID NO:10-18 were compared with sequences from the Incyte LIFESEQ database and public domain databases using BLAST and other implementations of the Smith-Waterman algorithm. Sequences from these databases that matched SEQ ID NO:10-18 were assembled into clusters of contiguous and overlapping sequences using assembly algorithms such as Phrap (Table 7). Radiation hybrid and genetic mapping data available from public resources such as the Stanford Human Genome Center (SHGC), Whitehead Institute for Genome Research (WIGR), and Généthon were used to determine if any of the clustered sequences had been previously mapped. Inclusion of a mapped sequence in a cluster resulted in the assignment of all sequences of that cluster, including its particular SEQ ID NO:, to that map location.

Map locations are represented by ranges, or intervals, of human chromosomes. The map position of an interval, in centiMorgans, is measured relative to the terminus of the chromosome's p-

arm. (The centiMorgan (cM) is a unit of measurement based on recombination frequencies between chromosomal markers. On average, 1 cM is roughly equivalent to 1 megabase (Mb) of DNA in humans, although this can vary widely due to hot and cold spots of recombination.) The cM distances are based on genetic markers mapped by Généthon which provide boundaries for radiation
 5 hybrid markers whose sequences were included in each of the clusters. Human genome maps and other resources available to the public, such as the NCBI "GeneMap'99" World Wide Web site (<http://www.ncbi.nlm.nih.gov/genemap/>), can be employed to determine if previously identified disease genes map within or in proximity to the intervals indicated above.

In this manner, SEQ ID NO:16 was mapped to chromosome 2 within the interval from
 10 190.80 to 197.60 centiMorgans.

VII. Analysis of Polynucleotide Expression

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel (1995)
 15 supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in cDNA databases such as GenBank or LIFESEQ (Incyte Genomics). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar.
 20 The basis of the search is the product score, which is defined as:

$$\frac{\text{BLAST Score} \times \text{Percent Identity}}{5 \times \text{minimum \{length(Seq. 1), length(Seq. 2)\}}}$$

25 The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. The product score is a normalized value between 0 and 100, and is calculated as follows: the BLAST score is multiplied by the percent nucleotide identity and the product is divided by (5 times the length of the shorter of the two sequences). The BLAST score is calculated by assigning a score of +5 for every base that matches in a high-scoring segment pair (HSP), and -4 for
 30 every mismatch. Two sequences may share more than one HSP (separated by gaps). If there is more than one HSP, then the pair with the highest BLAST score is used to calculate the product score. The product score represents a balance between fractional overlap and quality in a BLAST alignment. For example, a product score of 100 is produced only for 100% identity over the entire length of the shorter of the two sequences being compared. A product score of 70 is produced either by 100% identity and

70% overlap at one end, or by 88% identity and 100% overlap at the other. A product score of 50 is produced either by 100% identity and 50% overlap at one end, or 79% identity and 100% overlap.

Alternatively, polynucleotide sequences encoding XMES are analyzed with respect to the tissue sources from which they were derived. For example, some full length sequences are assembled, at least in part, with overlapping Incyte cDNA sequences (see Example III). Each cDNA sequence is derived from a cDNA library constructed from a human tissue. Each human tissue is classified into one of the following organ/tissue categories: cardiovascular system; connective tissue; digestive system; embryonic structures; endocrine system; exocrine glands; genitalia, female; genitalia, male; germ cells; hemic and immune system; liver; musculoskeletal system; nervous system; pancreas; respiratory system; sense organs; skin; stomatognathic system; unclassified/mixed; or urinary tract. The number of libraries in each category is counted and divided by the total number of libraries across all categories. Similarly, each human tissue is classified into one of the following disease/condition categories: cancer, cell line, developmental, inflammation, neurological, trauma, cardiovascular, pooled, and other, and the number of libraries in each category is counted and divided by the total number of libraries across all categories. The resulting percentages reflect the tissue- and disease-specific expression of cDNA encoding XMES. cDNA sequences and cDNA library/tissue information are found in the LIFESEQ GOLD database (Incyte Genomics, Palo Alto CA).

VIII. Extension of XMES Encoding Polynucleotides

Full length polynucleotide sequences were also produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer was synthesized to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and 2-mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C,

2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

5 The concentration of DNA in each well was determined by dispensing 100 µl PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 µl of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the
10 concentration of DNA. A 5 µl to 10 µl aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose gel to determine which reactions were successful in extending the sequence.

 The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For
15 shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on antibiotic-containing
20 media, and individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

 The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min; Step 5: steps 2, 3, and 4
25 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle
30 sequencing ready reaction kit (Applied Biosystems).

 In like manner, full length polynucleotide sequences are verified using the above procedure or are used to obtain 5' regulatory sequences using the above procedure along with oligonucleotides designed for such extension, and an appropriate genomic library.

IX. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:10-18 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ - 32 P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10^7 counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

X. Microarrays

The linkage or synthesis of array elements upon a microarray can be achieved utilizing photolithography, piezoelectric printing (ink-jet printing, See, e.g., Baldeschweiler, *supra*), mechanical microspotting technologies, and derivatives thereof. The substrate in each of the aforementioned technologies should be uniform and solid with a non-porous surface (Schena (1999), *supra*). Suggested substrates include silicon, silica, glass slides, glass chips, and silicon wafers. Alternatively, a procedure analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced using available methods and machines well known to those of ordinary skill in the art and may contain any appropriate number of elements. (See, e.g., Schena, M. et al. (1995) *Science* 270:467-470; Shalon, D. et al. (1996) *Genome Res.* 6:639-645; Marshall, A. and J. Hodgson (1998) *Nat. Biotechnol.* 16:27-31.)

Full length cDNAs, Expressed Sequence Tags (ESTs), or fragments or oligomers thereof may comprise the elements of the microarray. Fragments or oligomers suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). The array elements are hybridized with polynucleotides in a biological sample. The polynucleotides in the biological sample are conjugated to a fluorescent label or other molecular tag for ease of detection.

After hybridization, nonhybridized nucleotides from the biological sample are removed, and a fluorescence scanner is used to detect hybridization at each array element. Alternatively, laser desorption and mass spectrometry may be used for detection of hybridization. The degree of complementarity and the relative abundance of each polynucleotide which hybridizes to an element on the microarray may be assessed. In one embodiment, microarray preparation and usage is described in detail below.

Tissue or Cell Sample Preparation

Total RNA is isolated from tissue samples using the guanidinium thiocyanate method and poly(A)⁺ RNA is purified using the oligo-(dT) cellulose method. Each poly(A)⁺ RNA sample is reverse transcribed using MMLV reverse-transcriptase, 0.05 pg/ μ l oligo-(dT) primer (21mer), 1X first strand buffer, 0.03 units/ μ l RNase inhibitor, 500 μ M dATP, 500 μ M dGTP, 500 μ M dTTP, 40 μ M dCTP, 40 μ M dCTP-Cy3 (BDS) or dCTP-Cy5 (Amersham Pharmacia Biotech). The reverse transcription reaction is performed in a 25 ml volume containing 200 ng poly(A)⁺ RNA with GEMBRIGHT kits (Incyte). Specific control poly(A)⁺ RNAs are synthesized by in vitro transcription from non-coding yeast genomic DNA. After incubation at 37°C for 2 hr, each reaction sample (one with Cy3 and another with Cy5 labeling) is treated with 2.5 ml of 0.5M sodium hydroxide and incubated for 20 minutes at 85°C to stop the reaction and degrade the RNA. Samples are purified using two successive CHROMA SPIN 30 gel filtration spin columns (CLONTECH Laboratories, Inc. (CLONTECH), Palo Alto CA) and after combining, both reaction samples are ethanol precipitated using 1 ml of glycogen (1 mg/ml), 60 ml sodium acetate, and 300 ml of 100% ethanol. The sample is then dried to completion using a SpeedVAC (Savant Instruments Inc., Holbrook NY) and resuspended in 14 μ l 5X SSC/0.2% SDS.

Microarray Preparation

Sequences of the present invention are used to generate array elements. Each array element is amplified from bacterial cells containing vectors with cloned cDNA inserts. PCR amplification uses primers complementary to the vector sequences flanking the cDNA insert. Array elements are amplified in thirty cycles of PCR from an initial quantity of 1-2 ng to a final quantity greater than 5 μ g. Amplified array elements are then purified using SEPHACRYL-400 (Amersham Pharmacia Biotech).

Purified array elements are immobilized on polymer-coated glass slides. Glass microscope slides (Corning) are cleaned by ultrasound in 0.1% SDS and acetone, with extensive distilled water washes between and after treatments. Glass slides are etched in 4% hydrofluoric acid (VWR Scientific Products Corporation (VWR), West Chester PA), washed extensively in distilled water, and coated with 0.05% aminopropyl silane (Sigma) in 95% ethanol. Coated slides are cured in a 110°C oven.

Array elements are applied to the coated glass substrate using a procedure described in US Patent No. 5,807,522, incorporated herein by reference. 1 μ l of the array element DNA, at an average concentration of 100 ng/ μ l, is loaded into the open capillary printing element by a high-speed robotic apparatus. The apparatus then deposits about 5 nl of array element sample per slide.

- 5 Microarrays are UV-crosslinked using a STRATALINKER UV-crosslinker (Stratagene). Microarrays are washed at room temperature once in 0.2% SDS and three times in distilled water. Non-specific binding sites are blocked by incubation of microarrays in 0.2% casein in phosphate buffered saline (PBS) (Tropix, Inc., Bedford MA) for 30 minutes at 60°C followed by washes in 0.2% SDS and distilled water as before.

10 **Hybridization**

- Hybridization reactions contain 9 μ l of sample mixture consisting of 0.2 μ g each of Cy3 and Cy5 labeled cDNA synthesis products in 5X SSC, 0.2% SDS hybridization buffer. The sample mixture is heated to 65°C for 5 minutes and is aliquoted onto the microarray surface and covered with an 1.8 cm² coverslip. The arrays are transferred to a waterproof chamber having a cavity just slightly
15 larger than a microscope slide. The chamber is kept at 100% humidity internally by the addition of 140 μ l of 5X SSC in a corner of the chamber. The chamber containing the arrays is incubated for about 6.5 hours at 60°C. The arrays are washed for 10 min at 45°C in a first wash buffer (1X SSC, 0.1% SDS), three times for 10 minutes each at 45°C in a second wash buffer (0.1X SSC), and dried.

Detection

- 20 Reporter-labeled hybridization complexes are detected with a microscope equipped with an Innova 70 mixed gas 10 W laser (Coherent, Inc., Santa Clara CA) capable of generating spectral lines at 488 nm for excitation of Cy3 and at 632 nm for excitation of Cy5. The excitation laser light is focused on the array using a 20X microscope objective (Nikon, Inc., Melville NY). The slide containing the array is placed on a computer-controlled X-Y stage on the microscope and raster-
25 scanned past the objective. The 1.8 cm x 1.8 cm array used in the present example is scanned with a resolution of 20 micrometers.

- In two separate scans, a mixed gas multiline laser excites the two fluorophores sequentially. Emitted light is split, based on wavelength, into two photomultiplier tube detectors (PMT R1477, Hamamatsu Photonics Systems, Bridgewater NJ) corresponding to the two fluorophores. Appropriate
30 filters positioned between the array and the photomultiplier tubes are used to filter the signals. The emission maxima of the fluorophores used are 565 nm for Cy3 and 650 nm for Cy5. Each array is typically scanned twice, one scan per fluorophore using the appropriate filters at the laser source, although the apparatus is capable of recording the spectra from both fluorophores simultaneously.

- The sensitivity of the scans is typically calibrated using the signal intensity generated by a
35 cDNA control species added to the sample mixture at a known concentration. A specific location on

the array contains a complementary DNA sequence, allowing the intensity of the signal at that location to be correlated with a weight ratio of hybridizing species of 1:100,000. When two samples from different sources (e.g., representing test and control cells), each labeled with a different fluorophore, are hybridized to a single array for the purpose of identifying genes that are differentially expressed, the calibration is done by labeling samples of the calibrating cDNA with the two fluorophores and adding identical amounts of each to the hybridization mixture.

The output of the photomultiplier tube is digitized using a 12-bit RTI-835H analog-to-digital (A/D) conversion board (Analog Devices, Inc., Norwood MA) installed in an IBM-compatible PC computer. The digitized data are displayed as an image where the signal intensity is mapped using a linear 20-color transformation to a pseudocolor scale ranging from blue (low signal) to red (high signal). The data is also analyzed quantitatively. Where two different fluorophores are excited and measured simultaneously, the data are first corrected for optical crosstalk (due to overlapping emission spectra) between the fluorophores using each fluorophore's emission spectrum.

A grid is superimposed over the fluorescence signal image such that the signal from each spot is centered in each element of the grid. The fluorescence signal within each element is then integrated to obtain a numerical value corresponding to the average intensity of the signal. The software used for signal analysis is the GEMTOOLS gene expression analysis program (Incyte).

XI. Complementary Polynucleotides

Sequences complementary to the XMES-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring XMES. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of XMES. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the XMES-encoding transcript.

XII. Expression of XMES

Expression and purification of XMES is achieved using bacterial or virus-based expression systems. For expression of XMES in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac* (*tac*) hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3). Antibiotic resistant bacteria express XMES upon induction with isopropyl beta-D-

thiogalactopyranoside (IPTG). Expression of XMES in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding XMES by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

In most expression systems, XMES is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from XMES at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995, supra, ch. 10 and 16). Purified XMES obtained by these methods can be used directly in the assays shown in Examples XVI, and XVII where applicable.

XIII. Functional Assays

XMES function is assessed by expressing the sequences encoding XMES at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include PCMV SPORT (Life Technologies) and PCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-

based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of XMES on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding XMES and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding XMES and other genes of interest can be analyzed by northern analysis or microarray techniques.

XIV. Production of XMES Specific Antibodies

XMES substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the XMES amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Applied Biosystems) using FMOC chemistry and coupled to KLH (Sigma-Aldrich, St. Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-XMES activity by, for example, binding the peptide or XMES to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

XV. Purification of Naturally Occurring XMES Using Specific Antibodies

Naturally occurring or recombinant XMES is substantially purified by immunoaffinity chromatography using antibodies specific for XMES. An immunoaffinity column is constructed by covalently coupling anti-XMES antibody to an activated chromatographic resin, such as

- 5 CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is blocked and washed according to the manufacturer's instructions.

Media containing XMES are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of XMES (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt
10 antibody/XMES binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such as urea or thiocyanate ion), and XMES is collected.

XVI. Identification of Molecules Which Interact with XMES

XMES, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules
15 previously arrayed in the wells of a multi-well plate are incubated with the labeled XMES, washed, and any wells with labeled XMES complex are assayed. Data obtained using different concentrations of XMES are used to calculate values for the number, affinity, and association of XMES with the candidate molecules.

Alternatively, molecules interacting with XMES are analyzed using the yeast two-hybrid
20 system as described in Fields, S. and O. Song (1989) Nature 340:245-246, or using commercially available kits based on the two-hybrid system, such as the MATCHMAKER system (Clontech).

XMES may also be used in the PATHCALLING process (CuraGen Corp., New Haven CT) which employs the yeast two-hybrid system in a high-throughput manner to determine all interactions between the proteins encoded by two large libraries of genes (Nandabalan, K. et al. (2000) U.S. Patent
25 No. 6,057,101).

XVII. Demonstration of XMES Activity

An assay for XMES-1 activity measures its inhibitory activity on Hepatocyte Growth Factor (HGF) activator. In this assay, HGF activator (450 ng/ml) is mixed with various concentrations of purified XMES-1 in PBS containing 0.05% CHAPS and incubated at 37 degrees C for 30 minutes to
30 form an enzyme-inhibitor complex. The remaining HGF-converting activity in the mixture is measured by the addition of equal amounts of single chain HGF (sc-HGF) (1.5 µg/ml in PBS containing 0.05% CHAPS) and dextran sulfate (100 mg/ml, MWCO=500,000, Sigma) followed by further incubation for 2 hours, and subsequent analysis by SDS-PAGE under reducing gel conditions. The gel is stained with coomassie blue and the amounts of sc-HGF and the heterodimeric form are measured by scanning the

stained bands. The inhibitory activity of XMES-1 against HGF activator is estimated by calculating the ratio of the remaining single chain form to total HGF (Shimomura, T. et al. (1997) J. Biol. Chem. 272:6370-6376).

An assay for XMES activity measures cell proliferation as the amount of newly initiated DNA
5 synthesis in Swiss mouse 3T3 cells. A plasmid containing polynucleotides encoding XMES is
transfected into quiescent 3T3 cultured cells using methods well known in the art. The transiently
transfected cells are then incubated in the presence of [³H]thymidine, a radioactive DNA precursor.
Where applicable, varying amounts of XMES ligand are added to the transfected cells. Incorporation
of [³H]thymidine into acid-precipitable DNA is measured over an appropriate time interval, and the
10 amount incorporated is directly proportional to the amount of newly synthesized DNA.

Various modifications and variations of the described methods and systems of the invention will
be apparent to those skilled in the art without departing from the scope and spirit of the invention.
Although the invention has been described in connection with certain embodiments, it should be
understood that the invention as claimed should not be unduly limited to such specific embodiments.
15 Indeed, various modifications of the described modes for carrying out the invention which are obvious
to those skilled in molecular biology or related fields are intended to be within the scope of the following
claims.

Table 1

Incyte Project ID	Polypeptide SEQ ID NO:	Incyte Polypeptide ID	Polynucleotide SEQ ID NO:	Incyte Polynucleotide ID
1657368	1	1657368CD1	10	1657368CB1
4028972	2	4028972CD1	11	4028972CB1
5398353	3	5398353CD1	12	5398353CB1
71234118	4	71234118CD1	13	71234118CB1
240168	5	240168CD1	14	240168CB1
7481107	6	7481107CD1	15	7481107CB1
7476245	7	7476245CD1	16	7476245CB1
5819744	8	5819744CD1	17	5819744CB1
5378618	9	5378618CD1	18	5378618CB1

Table 2

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
1	165736CD1	g3721617	3.30E-42	[Mus musculus] mac25 insulin-like growth factor binding protein Komatsu S. et al. (2000) Biochem Biophys Res Commun 267:109-117
		g8217418	1.00E-169	[Homo sapiens] bA108L7.1 (novel Insulin-like growth factor binding type protein with Kazal-type serine protease inhibitor domain)
2	4028972CD1	g294600	0	[Rattus norvegicus] neurexin III-alpha Ushkaryov Y.A. and Sudhof T.C. (1993) Proc Natl Acad Sci U S A 1993 90:6410-6414
3	5398353CD1	g4514718	1.50E-22	FGF-19 [Homo sapiens] (Nishimura, T. (1999) Biochim. Biophys. Acta 1444:148-151)
		g9049445	1.00E-120	[Homo sapiens] FGF-21
4	71234118CD1	g11177164	4.00E-59	[Mus musculus] polydom protein
		g1438954	3.20E-18	[Homo sapiens] neuronal pentraxin 1
				Omeis, I.A. et al. (1996) Genomics 36:543-545
5	240168CD1	g2924601	1.10E-29	[Homo sapiens] hepatocyte growth factor activator inhibitor Shimomura, T. et al. (1997) J. Biol. Chem. 272:6370-6376

Table 2 (cont.)

Polypeptide SEQ ID NO:	Incyte Polypeptide ID	GenBank ID NO:	Probability score	GenBank Homolog
6	7481107CD1	g632490	7.50E-185	[Bos taurus] cartilage-derived morphogenetic protein Chang, S.C. et al. (1994) J. Biol. Chem. 269:28227-28234
7	7476245CD1	g189228	2.70E-54	[Homo sapiens] neuromedin B Krone, I.M. et al. (1988) J. Biol. Chem. 263:13317-13323
9	5378618CD1	g4590406	2.90E-19	[Drosophila melanogaster] slit protein Kidd, T. et al. (1999) Cell 96:785-794
		g9309467	0	[Mus musculus] leucine-rich glioma-inactivated 1 protein precursor

Table 3

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
1	1657368CD1	304	T250 Y251 T112 S117 T185	N159 N183 N277	M1-A30 Signal peptide Signal_cleavage: M1-A30 C53-C90 Insulin-like growth factor binding proteins V121-C168 Kazal-type serine protease inhibitor domain G186-G255 Immunoglobulin domain G50-D115 Insulin-like growth factor binding proteins signature L85-G171 Kazal serine protease inhibitors family signature V184-L271 MAC25 FOLLISTATINLIKE PROSTACYCLIN STIMULATING FACTOR	HMMER SPSCAN HMMER_PFAM HMMER_PFAM HMMER_PFAM HMMER_PFAM PROFILESKAN PROFILESKAN BLAST_PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
2	4028972CD1	1438	T11 S46 S100 T264 S290 T303 T469 S503 T531 T537 S564 S647 S714 S719 T730 T737 S762 T782 S791 T823 T878 S949 S972 T47 S79 S282 S283 T372 S392 T507 T708 T730 T795 T810 T906 Y273 Y547 Y770 Y860	N58 N105 N761 N403 N592 N776 N940 N943	C662-C673 Asx Hydroxyl S20-M31 Gataase_Type_I signal_cleavage: M1-G27 F10-G27, M1365-A1383 Transmembrane domain F55-G174, F291-R420, F479-N626, L740-I850, F903-D1034, F1126-W1201 Laminin G domain C202-C234, C651-C683, C1057-C1089 EGF-like domain S647-N658 Type II EGF-like signature E204-C224 COMPLEMENT C9 SIGNATURE F1098-Y1437 II-BETA NEUREXIN F717-E802, G803-Q902, G1210-T1338, T1338-V1438 ALTERNATIVE NEUREXIN PRECURSOR	MOTIFS MOTIFS SPSCAN HMMER HMMER_PFAM HMMER_PFAM BLIMPS_PRINTS BLIMPS_PRINTS BLAST_DOMO BLAST_PRODOM

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID.	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
3	5398353CD1	208	S121 T6 T97		signal_cleavage: M1-A27	SPSCAN
					Fibroblast growth factor FGF:H59-H139	HMMER_Pfam
					HEPARIN BINDING GROWTH FACTOR FAMILY PR00263B:L80-G94 PR00263C:F100-S112 PR00263D:P117-S136	BLIMPS_PRINTS
					IL1/HBGF FAMILY SIGNATURE PR00262A:P87-H114 PR00262B:A119-H139	BLIMPS_PRINTS
					HBGF/FGF family signature hbgef_fgf.prf:P87-G140	PROFILES SCAN
					GROWTH FACTOR FIBROBLAST MITOGEN SIGNAL HEPARIN BINDING VASCULARIZATION PD000831:H59-H139	BLAST_PRODOR
					HBGF/FGF FAMILY DM00642 P48803 47-205:S16-P165 DM00642 P48805 29-186:D32-P165 DM00642 P10767 50-207:G41-H139 DM00642 P05524 5-181:G40-P160	BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
4	71234118CD1	159	S114 S2 S69 Y75 N6 N85		signal_cleavage: M1-G35 Pentaxin family pentaxin: S80-K141 Pentaxin family proteins BL00289C:H122-G140 Pentaxin family signature pentaxin.prf:V100-A147 PENTAXIN SIGNATURE PR00895A:L58-D72 PR00895C:H122-G140 PRECURSOR SIGNAL PENTAXIN GLYCOPROTEIN PLASMA CREATINE CALCIUM ACUTE PHASE PD002153:Y48-G157 C-REACTIVE PROTEIN DM00835 P47971 194-431:F36-I155 DM00835 P47970 187-426:N37-I155	SPSCAN HMMER_PFAM BLIMPS_BLOCKS PROFILESCAN BLIMPS_PRINTS BLAST_PRODROM BLAST_DOMO

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
5	240168CD1	500	S11 S201 S234 S339 S34 S377 S382 S395 S422 S435 S53 T166 T197 T263 T279 T397 T487 Y431	N164 N291 N401	LDL RECEPTOR LIGAND-BINDING REPEAT DM00045 P98072 663-717:H314-L347 HEPATOCYTE GROWTH FACTOR ACTIVATOR INHIBITOR GLYCOPROTEIN PDI20361:G84-L297, S11-P44 LDL-receptor class A (LDLRA) domain proteins BL01209:C329-E341 LOW DENSITY LIPOPROTEIN PR00261:G320-E341 signal peptide signal peptide:M1-A37 transmembrane domain transmem domain:V452-C471 Low-density lipoprotein receptor domain ldl_recept_a:H308-N346 SpSCAN signal_cleavage:M1-A37	BLAST_DOMO BLAST_PRODUM BLIMPS_BLOCKS BLIMPS_PRINTS HMMER HMMER_PPFAM SPSCAN

Table 3 (cont.)

SEQ IID	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
6	7481107CD1	455	S106 S120 S129 S156 S284 S33 S355 S39 S393 S45 T3 T336	N114	TGF-BETA FAMILY DM00245 P43026 I74- 501-R331-R455, L99-G298 DM00245 P12644 95-408-R332- R455, A124-V217 DM00245 P12643 88-396-R345- R455, Q119-V217 DM00245 I49541 105- 420-R333-R455, A124-V217	BLAST_DOMO
					GLYCOPROTEIN PRECURSOR SIGNAL GROWTH FACTOR PROTEIN CYTOKINE BETA BONE MORPHOGENETIC PD000357:C354-R455 TGF-beta family proteins BL00250:C354-F389, C419- C454	BLAST_PRODOM BLIMPS_BLOCKS
					GROWTH FACTOR CYSTINE KNOT SUPERFAMILY SIGNATURE PR00438:E379-D388, E450- C454	BLIMPS_PRINTS
					INHIBIN ALPHA CHAIN SIGNATURE PR00669:C354-W371	HMMER
					signal peptide signal_peptide:M1-G22	HMMER_PFBAM
					Transforming growth factor beta like TGF-beta:C354- R455	
					TGF-beta propeptide TGFb_propeptide:G61-R268 TGF_beta:I372-C387	MOTIFS

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
7	7476245CD1	121	S76		BOMBESIN-LIKE PEPTIDES FAMILY DM00828 P08949 18-55:A18-M56 DM00828 P01297 1-31:A25-M56 NEUROMEDIN B32 PRECURSOR CONTAINS: B BOMBESIN FAMILY AMIDATION CLEAVAGE ON PAIR OF BASIC RESIDUES SIGNAL PD054439:G57-K121 NEUROMEDIN B32 CONTAINS: B BOMBESIN FAMILY AMIDATION PRECURSOR CLEAVAGE ON PD026110:A25-M56 Bombesin-like peptides family proteins BL00257:R46-M56 signal peptide signal_peptide:M1-D30 Bombesin:W50-M56 Bombesin-like peptides family signature bombesin.prf:D30-R81 SpSCAN signal_cleavage:M1-P26	BLAST_DOMO BLAST_PRODOM BLIMPS_BLOCKS HMMER MOTIFS PROFILESSCAN SPSCAN
8	5819744CD1	55	S12 S17 T2		Parathyroid hormone family signature parathyroid.prf:M1-K50 SpSCAN signal_cleavage:M1-P51	PROFILESSCAN SPSCAN

Table 3 (cont.)

SEQ ID NO:	Incyte Polypeptide ID	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequences, Domains and Motifs	Analytical Methods and Databases
9	5378618CD1	545	S154 S158 S238 S345 S366 S390 S404 S405 S425 S471 S528 T188 T209 T216 T374 T518 T72	N186 N271 N402 N70	Leucine-rich repeat signature PR00019:L156-F169, L135-L148 signal peptide Leucine Rich Repeat LR:D62-P85, S86-F109, H110-R133, D134-D157 Leucine rich repeat C-terminal domain LRRCT:N167-T216 Spscan signal_cleavage:M1-L23 Atp_Gtp_A binding site:A526-T533	BLIMPS_PRINTS HMMER HMMER_PFAM SPSCAN MOTIFS

Table 4

Poly-nucleotide SEQ ID NO:	Incyte Poly-nucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
10	1657368CB1	1374	1279-1374	7185657H1 (BONRREC01)	32	571
				1657373H1 (URETTTUT01)	1	227
				2687816F6 (LJUNGNOT23)	833	1374
				5851853H1 (FIBAUNT02)	684	942
				GNN.96967325_000005_006	199	1113
11	4028972CB1	4541	916-2416, 1-158, 2747-3709	2304926R6 (NGANNOT01)	4158	4541
				70032782D1	3771	4163
				70032296D1	3418	3776
				5498652F6 (BRABDIR01)	3659	4154
				70503374V1	516	1173
				70503718V1	2106	2675
				6608073H1 (BRAXDIC01)	3063	3519
				70505915V1	2705	3333
				70505907V1	1413	2072
				6993321H1 (BRAQTDRO2)	1	561
				70501232V1	784	1409
				70504786V1	1385	2066
12	5398353CB1	1117	485-916, 1-75, 990-1117, 157-211	70504213V1	1997	2655
				70505588V1	2429	3132
				71281268V1	1	631
				71152050V1	682	1117
				71154676V1	657	1114
13	71234118CB1	2460	2070-2175, 1-38, 1692-1726, 1395-1534, 156-1013, 2409-2460	5398353T6 (LIVRTUT13)	506	1084
				4062433T6 (BRAINT021)	1798	2460
				71008692V1	219	836
				71234448V1	1	590
				71010865V1	1150	1746
				5647010F8 (BRAITUT23)	1727	2360
				71009374V1	978	1560
				71008960V1	599	1085

Table 4 (cont.)

Poly-nucleotide SEQ ID NO:	Incyte Poly-nucleotide ID	Sequence Length	Selected Fragment(s)	Sequence Fragments	5' Position	3' Position
14	240168CB1	2601	592-648, 1-45, 706-754	3227038H1 (EPIGNOT01) g3153915 1833192H1 (BRAINON01) 1750805F6 (LIVRUT01) 6764130J1 (BRAUNOR01) 1913790H1 (PROSTUT04) 6975777H1 (BRAHTR04) 7070876H1 (BRAUTDR02) 7653484H1 (UTREDME06) GNN.g7768040_000033_002	1348 649 2367 2189 657 2340 1357 1058 1 147	1626 1174 2601 2580 1344 2587 2035 1545 471 759
15	7481107CB1	2791	1-1312, 2425-2791, 1841-1883, 1598-1658	1451206F6 (PENITUT01) GNN.g8439934_000008_002. edit.1 1387140F6 (CARGDIT02) 6152346H1 (ENDMONT04) GBL.chr8_smoosh_4.comp 2377775F6 (ISLITNOT01) 1577577F6 (LNODNOT03) 5813552F8 (PROSTUS23) 5813552F8 (PROSTUS23)	1618 1029 2188 2165 1 411 1 210 1	2210 2396 2791 2451 1427 709 554 753 636
16	7476245CB1	709	651-709	6718171H1 (CONDUT02) 70656442V1	1805 2901	2358 3413
17	5819744CB1	753	1-753	GNN.g4416547_000001_4 8264863U1 FL5378618_g4416547_g4091 819	78 1683 78	1715 2349 1715
18	5378618CB1	3413	3392-3413, 1-1039, 1696-2626	GBI.g4416547.fasta.edit 70657537V1	1 2340	201 2945

Table 5

Polynucleotide SEQ ID NO:	Incyte Project ID	Representative Library
10	1657368CB1	LUNGNOT23
11	4028972CB1	BRADIR01
12	5398353CB1	LIVRTUT13
13	71234118CB1	PLACNOB01
14	240168CB1	LIVRTUT01
15	7481107CB1	CARGIT02
16	7476245CB1	BRAITUT12
17	5819744CB1	PROSTUS23
18	5378618CB1	BRSTTUT03

Table 6

Library	Vector	Library Description
BRABDIR01	pINCY	Library was constructed using RNA isolated from diseased cerebellum tissue removed from the brain of a 57-year-old Caucasian male, who died from a cerebrovascular accident. Patient history included Huntington's disease, emphysema, and tobacco abuse.
BRAITUT12	pINCY	Library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 40-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated grade 4 gemistocytic astrocytoma.
BRSTTUT03	PSPORT1	Library was constructed using RNA isolated from breast tumor tissue removed from a 58-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated multicentric invasive grade 4 lobular carcinoma. The mass was identified in the upper outer quadrant, and three separate nodules were found in the lower outer quadrant of the left breast. Patient history included skin cancer, rheumatic heart disease, osteoarthritis, and tuberculosis. Family history included cerebrovascular disease, coronary artery aneurysm, breast cancer, prostate cancer, atherosclerotic coronary artery disease, and type I diabetes.
CARGDIT02	pINCY	Library was constructed using RNA isolated from cartilage obtained from 4 donors with end-stage osteoarthritis. The patients had received a variety of non-steroidal anti-inflammatory drugs.
LIVRTUT01	pINCY	Library was constructed using RNA isolated from liver tumor tissue removed from a 51-year-old Caucasian female during a hepatic lobectomy. Pathology indicated metastatic grade 3 adenocarcinoma consistent with colon cancer. Family history included a malignant neoplasm of the liver.

Table 6 (cont.)

Library	Vector	Library Description
LIVRTUT13	pINCY	Library was constructed using RNA isolated from liver tumor tissue removed from a 62-year-old Caucasian female during partial hepatectomy and exploratory laparotomy. Pathology indicated metastatic intermediate grade neuroendocrine carcinoma, consistent with islet cell tumor, forming nodules ranging in size, in the lateral and medial left liver lobe. The pancreas showed fibrosis, chronic inflammation and fat necrosis consistent with pseudocyst. The gall bladder showed mild chronic cholecystitis. Patient history included malignant neoplasm of the pancreas tail, pulmonary embolism, hyperlipidemia, thrombophlebitis, joint pain in multiple joints, type II diabetes, benign hypertension, and cerebrovascular disease. Family history included pancreas cancer, secondary liver cancer, benign hypertension, and hyperlipidemia.
LUNGNOT23	pINCY	Library was constructed using RNA isolated from left lobe lung tissue removed from a 58-year-old Caucasian male. Pathology for the associated tumor tissue indicated metastatic grade 3 (of 4) osteosarcoma. Patient history included soft tissue cancer, secondary cancer of the lung, prostate cancer, and an acute duodenal ulcer with hemorrhage. Family history included prostate cancer, breast cancer, and acute leukemia.
PLACNOB01	PBLUESCRIPT	Library was constructed using RNA isolated from placenta.

Table 6 (cont.)

Library	Vector	Library Description
PROTUS23	pINCY	<p>This subtracted prostate tumor library was constructed using 10 million clones from a pooled prostate tumor library that was subjected to 2 rounds of subtractive hybridization with 10 million clones from a pooled prostate tissue library. The starting library for subtraction was constructed by pooling equal numbers of clones from 4 prostate tumor libraries using mRNA isolated from prostate tumor removed from Caucasian males at ages 58 (A), 61 (B), 66 (C), and 68 (D) during prostatectomy with lymph node excision. Pathology indicated adenocarcinoma in all donors. History included elevated PSA, induration and tobacco abuse in donor A; elevated PSA, induration, prostate hyperplasia, renal failure, osteoarthritis, renal artery stenosis, benign HTN, thrombocytopenia, hyperlipidemia, tobacco/alcohol abuse and hepatitis C (carrier) in donor B; elevated PSA, induration, and tobacco abuse in donor C; and elevated PSA, induration, hypercholesterolemia, and kidney calculus in donor D. The hybridization probe for subtraction was constructed by pooling equal numbers of cDNA clones from 3 prostate tissue libraries derived from prostate tissue, prostate epithelial cells, and fibroblasts from prostate stroma from 3 different donors. Subtractive hybridization conditions were based on the methodologies of Swaroop et al., NAR 19 (1991):1954 and Bonaldo, et al. Genome Research 6 (1996):791.</p>

Table 7

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410; Altschul, S.F. et al. (1997) Nucleic Acids Res. 25:3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) Proc. Natl. Acad. Sci. USA 85:2444-2448; Pearson, W.R. (1990) Methods Enzymol. 183:63-98; and Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater; fastx E value=1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A Blocks IMProved Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff (1991) Nucleic Acids Res. 19:6565-6572; Henikoff, J.G. and S. Henikoff (1996) Methods Enzymol. 266:88-105; and Attwood, T.K. et al. (1997) J. Chem. Inf. Comput. Sci. 37:417-424.	Probability value= 1.0E-3 or less
HMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) J. Mol. Biol. 235:1501-1531; Sonnhammer, E.L.L. et al. (1998) Nucleic Acids Res. 26:320-322; Durbin, R. et al. (1998) Our World View, in a Nutshell, Cambridge Univ. Press, pp. 1-350.	PFAM hits: Probability value= 1.0E-3 or less Signal peptide hits: Score= 0 or greater

Table 7 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, M. et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221.	Normalized quality score>GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M.S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M.S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies.	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPSscan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12:431-439.	Score=3.5 or greater
TMAP	A program that uses weight matrices to delineate transmembrane segments on protein sequences and determine orientation.	Persson, B. and P. Argos (1994) J. Mol. Biol. 237:182-192; Persson, B. and P. Argos (1996) Protein Sci. 5:363-371.	
TMHMMER	A program that uses a hidden Markov model (HMM) to delineate transmembrane segments on protein sequences and determine orientation.	Sonnhammer, E.L. et al. (1998) Proc. Sixth Intl. Conf. on Intelligent Systems for Mol. Biol., Glasgow et al., eds., The Am. Assoc. for Artificial Intelligence Press, Menlo Park, CA, pp. 175-182.	
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch, A. et al. (1997) Nucleic Acids Res. 25:217-221; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide selected from the group consisting of:
 - a) a polypeptide comprising an amino acid sequence selected from the group consisting of
 - 5 SEQ ID NO:1-9,
 - b) a naturally occurring polypeptide comprising an amino acid sequence at least 90% identical to an amino acid sequence selected from the group consisting of SEQ ID NO:1-9,
 - c) a biologically active fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, and
 - 10 d) an immunogenic fragment of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9.
2. An isolated polypeptide of claim 1 selected from the group consisting of SEQ ID NO:1-9.
- 15 3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide encoding a polypeptide of claim 2.
5. An isolated polynucleotide of claim 4 selected from the group consisting of SEQ ID
- 20 NO:10-18.
6. A recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide of claim 3.
- 25 7. A cell transformed with a recombinant polynucleotide of claim 6.
8. A transgenic organism comprising a recombinant polynucleotide of claim 6.
9. A method for producing a polypeptide of claim 1, the method comprising:
 - 30 a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and
 - b) recovering the polypeptide so expressed.
- 35

10. An isolated antibody which specifically binds to a polypeptide of claim 1.
11. An isolated polynucleotide selected from the group consisting of:
- a) a polynucleotide comprising a polynucleotide sequence selected from the group consisting
 - 5 of SEQ ID NO:10-18,
 - b) a naturally occurring polynucleotide comprising a polynucleotide sequence at least 90% identical to a polynucleotide sequence selected from the group consisting of SEQ ID NO:10-18,
 - c) a polynucleotide complementary to a polynucleotide of a),
 - d) a polynucleotide complementary to a polynucleotide of b), and
 - 10 e) an RNA equivalent of a)-d).
12. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 11.
13. A method for detecting a target polynucleotide in a sample, said target polynucleotide
- 15 having a sequence of a polynucleotide of claim 11, the method comprising:
- a) hybridizing the sample with a probe comprising at least 20 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization
 - 20 complex is formed between said probe and said target polynucleotide or fragments thereof, and
 - b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.
14. A method of claim 13, wherein the probe comprises at least 60 contiguous nucleotides.
- 25 15. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 11, the method comprising:
- a) amplifying said target polynucleotide or fragment thereof using polymerase chain reaction amplification, and
 - 30 b) detecting the presence or absence of said amplified target polynucleotide or fragment thereof, and, optionally, if present, the amount thereof.
16. A composition comprising a polypeptide of claim 1 and a pharmaceutically acceptable excipient.
- 35

17. A composition of claim 16, wherein the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NO:1-9.

18. A method for treating a disease or condition associated with decreased expression of functional XMES, comprising administering to a patient in need of such treatment the composition of claim 16.

19. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

20. A composition comprising an agonist compound identified by a method of claim 19 and a pharmaceutically acceptable excipient.

21. A method for treating a disease or condition associated with decreased expression of functional XMES, comprising administering to a patient in need of such treatment a composition of claim 20.

22. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

23. A composition comprising an antagonist compound identified by a method of claim 22 and a pharmaceutically acceptable excipient.

24. A method for treating a disease or condition associated with overexpression of functional XMES, comprising administering to a patient in need of such treatment a composition of claim 23.

25. A method of screening for a compound that specifically binds to the polypeptide of claim 1, said method comprising the steps of:

- a) combining the polypeptide of claim 1 with at least one test compound under suitable conditions, and

b) detecting binding of the polypeptide of claim 1 to the test compound, thereby identifying a compound that specifically binds to the polypeptide of claim 1.

26. A method of screening for a compound that modulates the activity of the polypeptide of claim 1, said method comprising:

- a) combining the polypeptide of claim 1 with at least one test compound under conditions permissive for the activity of the polypeptide of claim 1,
- b) assessing the activity of the polypeptide of claim 1 in the presence of the test compound, and
- c) comparing the activity of the polypeptide of claim 1 in the presence of the test compound with the activity of the polypeptide of claim 1 in the absence of the test compound, wherein a change in the activity of the polypeptide of claim 1 in the presence of the test compound is indicative of a compound that modulates the activity of the polypeptide of claim 1.

27. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 5, the method comprising:

- a) exposing a sample comprising the target polynucleotide to a compound, under conditions suitable for the expression of the target polynucleotide,
- b) detecting altered expression of the target polynucleotide, and
- c) comparing the expression of the target polynucleotide in the presence of varying amounts of the compound and in the absence of the compound.

28. A method for assessing toxicity of a test compound, said method comprising:

- a) treating a biological sample containing nucleic acids with the test compound;
- b) hybridizing the nucleic acids of the treated biological sample with a probe comprising at least 20 contiguous nucleotides of a polynucleotide of claim 11 under conditions whereby a specific hybridization complex is formed between said probe and a target polynucleotide in the biological sample, said target polynucleotide comprising a polynucleotide sequence of a polynucleotide of claim 11 or fragment thereof;
- c) quantifying the amount of hybridization complex; and
- d) comparing the amount of hybridization complex in the treated biological sample with the amount of hybridization complex in an untreated biological sample, wherein a difference in the amount of hybridization complex in the treated biological sample is indicative of toxicity of the test compound.

29. A diagnostic test for a condition or disease associated with the expression of XMES in a biological sample comprising the steps of:

- 5 a) combining the biological sample with an antibody of claim 10, under conditions suitable for the antibody to bind the polypeptide and form an antibody:polypeptide complex; and
- b) detecting the complex, wherein the presence of the complex correlates with the presence of the polypeptide in the biological sample.

30. The antibody of claim 10, wherein the antibody is:

- 10 a) a chimeric antibody,
- b) a single chain antibody,
- c) a Fab fragment,
- d) a F(ab')₂ fragment, or
- 15 e) a humanized antibody.

31. A composition comprising an antibody of claim 10 and an acceptable excipient.

32. A method of diagnosing a condition or disease associated with the expression of XMES in a subject, comprising administering to said subject an effective amount of the composition of claim 20 31.

33. A composition of claim 31, wherein the antibody is labeled.

34. A method of diagnosing a condition or disease associated with the expression of XMES in a subject, comprising administering to said subject an effective amount of the composition of claim 25 33.

35. A method of preparing a polyclonal antibody with the specificity of the antibody of claim 10 comprising:

- 30 a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, or an immunogenic fragment thereof, under conditions to elicit an antibody response;
- b) isolating antibodies from said animal; and

c) screening the isolated antibodies with the polypeptide, thereby identifying a polyclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9.

5 36. An antibody produced by a method of claim 35.

37. A composition comprising the antibody of claim 36 and a suitable carrier.

10 38. A method of making a monoclonal antibody with the specificity of the antibody of claim 10 comprising:

a) immunizing an animal with a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9, or an immunogenic fragment thereof, under conditions to elicit an antibody response;

b) isolating antibody producing cells from the animal;

15 c) fusing the antibody producing cells with immortalized cells to form monoclonal antibody-producing hybridoma cells;

d) culturing the hybridoma cells; and

e) isolating from the culture monoclonal antibody which binds specifically to a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9.

20

39. A monoclonal antibody produced by a method of claim 38.

40. A composition comprising the antibody of claim 39 and a suitable carrier.

25 41. The antibody of claim 10, wherein the antibody is produced by screening a Fab expression library.

42. The antibody of claim 10, wherein the antibody is produced by screening a recombinant immunoglobulin library.

30

43. A method for detecting a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9 in a sample, comprising the steps of:

a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and

b) detecting specific binding, wherein specific binding indicates the presence of a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9 in the sample.

5 44. A method of purifying a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9 from a sample, the method comprising:

a) incubating the antibody of claim 10 with a sample under conditions to allow specific binding of the antibody and the polypeptide; and

b) separating the antibody from the sample and obtaining the purified polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NO:1-9.

45. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:1.

46. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:2.

15

47. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:3.

48. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:4.

20

49. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:5.

50. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:6.

51. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:7.

25

52. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:8.

53. A polypeptide of claim 1, comprising the amino acid sequence of SEQ ID NO:9.

30

54. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:10.

55. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:11.

56. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:12.

57. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
5 NO:13.

58. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID NO:14.

59. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
10 NO:15.

60. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
NO:16.

61. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
15 NO:17.

62. A polynucleotide of claim 11, comprising the polynucleotide sequence of SEQ ID
20 NO:18.

<110> INCYTE GENOMICS, INC.

LAL, Preeti
YUE, Henry
HE, Ann
NGUYEN, Danniel B.
WALIA, Narinder
GANDHI, Aameena R.
AZIMZAI, Yalda
BANDMAN, Olga
TANG, Y. Tom
LU, Yan
BAUGHN, Mariah R.
DUGGAN, Brendan M.
LEE, Sally
HAFALIA, April
POLICKY, Jennifer L.

<120> EXTRACELLULAR MESSENGERS

<130> PF-0780 PCT.

<140> To Be Assigned

<141> Herewith

<150> 60/210,233; 60/213,465; 60/249,019

<151> 2000-06-06; 2000-06-23; 2000-11-14

<160> 18

<170> PERL Program

<210> 1

<211> 304

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 1657368CD1

<400> 1

Met	Leu	Pro	Pro	Pro	Arg	Pro	Ala	Ala	Ala	Leu	Ala	Leu	Pro	Val
1				5					10					15
Leu	Leu	Leu	Leu	Leu	Val	Val	Leu	Thr	Pro	Pro	Pro	Thr	Gly	Ala
				20					25					30
Arg	Pro	Ser	Pro	Gly	Pro	Asp	Tyr	Leu	Arg	Arg	Gly	Trp	Met	Arg
				35					40					45
Leu	Leu	Ala	Glu	Gly	Glu	Gly	Cys	Ala	Pro	Cys	Arg	Pro	Glu	Glu
				50					55					60
Cys	Ala	Ala	Pro	Arg	Gly	Cys	Leu	Ala	Gly	Arg	Val	Arg	Asp	Ala
				65					70					75
Cys	Gly	Cys	Cys	Trp	Glu	Cys	Ala	Asn	Leu	Glu	Gly	Gln	Leu	Cys
				80					85					90
Asp	Leu	Asp	Pro	Ser	Ala	His	Phe	Tyr	Gly	His	Cys	Gly	Glu	Gln
				95					100					105
Leu	Glu	Cys	Arg	Leu	Asp	Thr	Gly	Gly	Asp	Leu	Ser	Arg	Gly	Glu
				110					115					120
Val	Pro	Glu	Pro	Leu	Cys	Ala	Cys	Arg	Ser	Gln	Ser	Pro	Leu	Cys
				125					130					135
Gly	Ser	Asp	Gly	His	Thr	Tyr	Ser	Gln	Ile	Cys	Arg	Leu	Gln	Glu
				140					145					150
Ala	Ala	Arg	Ala	Arg	Pro	Asp	Ala	Asn	Leu	Thr	Val	Ala	His	Pro
				155					160					165
Gly	Pro	Cys	Glu	Ser	Gly	Pro	Gln	Ile	Val	Ser	His	Pro	Tyr	Asp
				170					175					180
Thr	Trp	Asn	Val	Thr	Gly	Gln	Asp	Val	Ile	Phe	Gly	Cys	Glu	Val
				185					190					195
Phe	Ala	Tyr	Pro	Met	Ala	Ser	Ile	Glu	Trp	Arg	Lys	Asp	Gly	Leu

Asp	Ile	Gln	Leu	200	Pro	Gly	Asp	Asp	Pro	205	His	Ile	Ser	Val	Gln	210
				215						220						225
Arg	Gly	Gly	Pro	230	Gln	Arg	Phe	Glu	Val	235	Thr	Gly	Trp	Leu	Gln	240
Gln	Ala	Val	Arg	245	Pro	Ser	Asp	Glu	Gly	250	Thr	Tyr	Arg	Cys	Leu	255
Arg	Asn	Ala	Leu	260	Gly	Gln	Val	Glu	Ala	265	Pro	Ala	Ser	Leu	Thr	270
Leu	Thr	Pro	Asp	275	Gln	Leu	Asn	Ser	Thr	280	Gly	Ile	Pro	Gln	Leu	285
Ser	Leu	Asn	Leu	290	Val	Pro	Glu	Glu	Glu	295	Ala	Glu	Ser	Glu	Glu	300
Asp	Asp	Tyr	Tyr													

<210> 2
 <211> 1438
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 4028972CD1

<400> 2

Met	Ser	Ser	Thr	Leu	His	Ser	Val	Phe	Phe	Thr	Leu	Lys	Val	Ser
1				5					10					15
Ile	Leu	Leu	Gly	Ser	Leu	Leu	Gly	Leu	Cys	Leu	Gly	Leu	Glu	Phe
				20					25					30
Met	Gly	Leu	Pro	Asn	Gln	Trp	Ala	Arg	Tyr	Leu	Arg	Trp	Asp	Ala
				35					40					45
Ser	Thr	Arg	Ser	Asp	Leu	Ser	Phe	Gln	Phe	Lys	Thr	Asn	Val	Ser
				50					55					60
Thr	Gly	Leu	Leu	Leu	Tyr	Leu	Asp	Asp	Gly	Gly	Val	Cys	Asp	Phe
				65					70					75
Leu	Cys	Leu	Ser	Leu	Val	Asp	Gly	Arg	Val	Gln	Leu	Arg	Phe	Ser
				80					85					90
Met	Asp	Cys	Ala	Glu	Thr	Ala	Val	Leu	Ser	Asn	Lys	Gln	Val	Asn
				95					100					105
Asp	Ser	Ser	Trp	His	Phe	Leu	Met	Val	Ser	Arg	Asp	Arg	Leu	Arg
				110					115					120
Thr	Val	Leu	Met	Leu	Asp	Gly	Glu	Gly	Gln	Ser	Gly	Glu	Leu	Gln
				125					130					135
Pro	Gln	Arg	Pro	Tyr	Met	Asp	Val	Val	Ser	Asp	Leu	Phe	Leu	Gly
				140					145					150
Gly	Val	Pro	Thr	Asp	Ile	Arg	Pro	Ser	Ala	Leu	Thr	Leu	Asp	Gly
				155					160					165
Val	Gln	Ala	Met	Pro	Gly	Phe	Lys	Gly	Leu	Ile	Leu	Asp	Leu	Lys
				170					175					180
Tyr	Gly	Asn	Ser	Glu	Pro	Arg	Leu	Leu	Gly	Ser	Arg	Gly	Val	Gln
				185					190					195
Met	Asp	Ala	Glu	Gly	Pro	Cys	Gly	Glu	Arg	Pro	Cys	Glu	Asn	Gly
				200					205					210
Gly	Ile	Cys	Phe	Leu	Leu	Asp	Gly	His	Pro	Thr	Cys	Asp	Cys	Ser
				215					220					225
Thr	Thr	Gly	Tyr	Gly	Gly	Lys	Leu	Cys	Ser	Glu	Asp	Val	Ser	Gln
				230					235					240
Asp	Pro	Gly	Leu	Ser	His	Leu	Met	Met	Ser	Glu	Gln	Gly	Arg	Ser
				245					250					255
Lys	Ala	Arg	Glu	Glu	Asn	Val	Ala	Thr	Phe	Arg	Gly	Ser	Glu	Tyr
				260					265					270
Leu	Cys	Tyr	Asp	Leu	Ser	Gln	Asn	Pro	Ile	Gln	Ser	Ser	Ser	Asp
				275					280					285
Glu	Ile	Thr	Leu	Ser	Phe	Lys	Thr	Trp	Gln	Arg	Asn	Gly	Leu	Ile
				290					295					300
Leu	His	Thr	Gly	Lys	Ser	Ala	Asp	Tyr	Val	Asn	Leu	Ala	Leu	Lys

Asp Gly Ala Val	305	Ser Leu Val Ile Asn	310	Leu Gly Ser Gly Ala	315
	320		325		330
Glu Ala Ile Val	335	Glu Pro Val Asn Gly	340	Lys Phe Asn Asp Asn	345
Trp His Asp Val	350	Lys Val Thr Arg Asn	355	Leu Arg Gln Val Thr	360
Ser Val Asp Gly	365	Ile Leu Thr Thr Thr	370	Gly Tyr Thr Gln Glu	375
Tyr Thr Met Leu	380	Gly Ser Asp Asp Phe	385	Phe Tyr Val Gly Gly	390
Pro Ser Thr Ala	395	Asp Leu Pro Gly Ser	400	Pro Val Ser Asn Asn	405
Met Gly Cys Leu	410	Lys Glu Val Val Tyr	415	Lys Asn Asn Asp Ile	420
Leu Glu Leu Ser	425	Arg Leu Ala Arg Ile	430	Ala Asp Thr Lys Met	435
Ile Tyr Gly Glu	440	Val Val Phe Lys Cys	445	Glu Asn Val Ala Thr	450
Asp Pro Ile Asn	455	Phe Glu Thr Pro Glu	460	Ala Tyr Ile Ser Leu	465
Lys Trp Asn Thr	470	Lys Arg Met Gly Ser	475	Ile Ser Phe Asp Phe	480
Thr Thr Glu Pro	485	Asn Gly Leu Ile Leu	490	Phe Thr His Gly Lys	495
Gln Glu Arg Lys	500	Asp Ala Arg Ser Gln	505	Lys Asn Thr Lys Val	510
Phe Phe Ala Val	515	Glu Leu Leu Asp Gly	520	Asn Leu Tyr Leu Leu	525
Asp Met Gly Ser	530	Gly Thr Ile Lys Val	535	Lys Ala Thr Gln Lys	540
Ala Asn Asp Gly	545	Glu Trp Tyr His Val	550	Asp Ile Gln Arg Asp	555
Arg Ser Gly Thr	560	Ile Ser Val Asn Ser	565	Arg Arg Thr Pro Phe	570
Ala Ser Gly Glu	575	Ser Glu Ile Leu Asp	580	Leu Glu Gly Asp Met	585
Leu Gly Gly Leu	590	Pro Glu Asn Arg Ala	595	Gly Leu Ile Leu Pro	600
Glu Leu Trp Thr	605	Ala Met Leu Asn Tyr	610	Gly Tyr Val Gly Cys	615
Arg Asp Leu Phe	620	Ile Asp Gly Arg Ser	625	Lys Asn Ile Arg Gln	630
Ala Glu Met Gln	635	Asn Ala Ala Gly Val	640	Lys Ser Ser Cys Ser	645
Met Ser Ala Lys	650	Gln Cys Asp Ser Tyr	655	Pro Cys Lys Asn Asn	660
Val Cys Lys Asp	665	Gly Trp Asn Arg Phe	670	Ile Cys Asp Cys Thr	675
Thr Gly Tyr Trp	680	Gly Arg Thr Cys Glu	685	Arg Glu Ala Ser Ile	690
Ser Tyr Asp Gly	695	Ser Met Tyr Met Lys	700	Ile Ile Met Pro Met	705
Met His Thr Glu	710	Ala Glu Asp Val Ser	715	Phe Arg Phe Met Ser	720
Arg Ala Tyr Gly	725	Leu Leu Val Ala Thr	730	Thr Ser Arg Asp Ser	735
Asp Thr Leu Arg	740	Leu Glu Leu Asp Gly	745	Gly Arg Val Lys Leu	750
Val Asn Leu Asp	755	Cys Ile Arg Ile Asn	760	Cys Asn Ser Ser Lys	765
Pro Glu Thr Leu	770	Tyr Ala Gly Gln Lys	775	Leu Asn Asp Asn Glu	780
His Thr Val Arg	785	Val Val Arg Arg Gly	790	Lys Ser Leu Lys Leu	795
Val Asp Asp Asp	800	Val Ala Glu Gly Thr	805	Met Val Gly Asp His	810

Arg	Leu	Glu	Phe	His	Asn	Ile	Glu	Thr	Gly	Ile	Met	Thr	Glu	Lys
				815					820					825
Arg	Tyr	Ile	Ser	Val	Val	Pro	Ser	Ser	Phe	Ile	Gly	His	Leu	Gln
				830					835					840
Ser	Leu	Met	Phe	Asn	Gly	Leu	Leu	Tyr	Ile	Asp	Leu	Cys	Lys	Asn
				845					850					855
Gly	Asp	Ile	Asp	Tyr	Cys	Glu	Leu	Lys	Ala	Arg	Phe	Gly	Leu	Arg
				860					865					870
Asn	Ile	Ile	Ala	Asp	Pro	Val	Thr	Phe	Lys	Thr	Lys	Ser	Ser	Tyr
				875					880					885
Leu	Ser	Leu	Ala	Thr	Leu	Gln	Ala	Tyr	Thr	Ser	Met	His	Leu	Phe
				890					895					900
Phe	Gln	Phe	Lys	Thr	Thr	Ser	Pro	Asp	Gly	Phe	Ile	Leu	Phe	Asn
				905					910					915
Ser	Gly	Asp	Gly	Asn	Asp	Phe	Ile	Ala	Val	Glu	Leu	Val	Lys	Gly
				920					925					930
Tyr	Ile	His	Tyr	Val	Phe	Asp	Leu	Gly	Asn	Gly	Pro	Asn	Val	Ile
				935					940					945
Lys	Gly	Asn	Ser	Asp	Arg	Pro	Leu	Asn	Asp	Asn	Gln	Trp	His	Asn
				950					955					960
Val	Val	Ile	Thr	Arg	Asp	Asn	Ser	Asn	Thr	His	Ser	Leu	Lys	Val
				965					970					975
Asp	Thr	Lys	Val	Val	Thr	Gln	Val	Ile	Asn	Gly	Ala	Lys	Asn	Leu
				980					985					990
Asp	Leu	Lys	Gly	Asp	Leu	Tyr	Met	Ala	Gly	Leu	Ala	Gln	Gly	Met
				995					1000					1005
Tyr	Ser	Asn	Leu	Pro	Lys	Leu	Val	Ala	Ser	Arg	Asp	Gly	Phe	Gln
				1010					1015					1020
Gly	Cys	Leu	Ala	Ser	Val	Asp	Leu	Asn	Gly	Arg	Leu	Pro	Asp	Leu
				1025					1030					1035
Ile	Asn	Asp	Ala	Leu	His	Arg	Ser	Gly	Gln	Ile	Glu	Arg	Gly	Cys
				1040					1045					1050
Glu	Gly	Pro	Ser	Thr	Thr	Cys	Gln	Glu	Asp	Ser	Cys	Ala	Asn	Gln
				1055					1060					1065
Gly	Val	Cys	Met	Gln	Gln	Trp	Glu	Gly	Phe	Thr	Cys	Asp	Cys	Ser
				1070					1075					1080
Met	Thr	Ser	Tyr	Ser	Gly	Asn	Gln	Cys	Asn	Asp	Pro	Gly	Ala	Thr
				1085					1090					1095
Tyr	Ile	Phe	Gly	Lys	Ser	Gly	Gly	Leu	Ile	Leu	Tyr	Thr	Trp	Pro
				1100					1105					1110
Ala	Asn	Asp	Arg	Pro	Ser	Thr	Arg	Ser	Asp	Arg	Leu	Ala	Val	Gly
				1115					1120					1125
Phe	Ser	Thr	Thr	Val	Lys	Asp	Gly	Ile	Leu	Val	Arg	Ile	Asp	Ser
				1130					1135					1140
Ala	Pro	Gly	Leu	Gly	Asp	Phe	Leu	Gln	Leu	His	Ile	Glu	Gln	Gly
				1145					1150					1155
Lys	Ile	Gly	Val	Val	Phe	Asn	Ile	Gly	Thr	Val	Asp	Ile	Ser	Ile
				1160					1165					1170
Lys	Glu	Glu	Arg	Thr	Pro	Val	Asn	Asp	Gly	Lys	Tyr	His	Val	Val
				1175					1180					1185
Arg	Phe	Thr	Arg	Asn	Gly	Gly	Asn	Ala	Thr	Leu	Gln	Val	Asp	Asn
				1190					1195					1200
Trp	Pro	Val	Asn	Glu	His	Tyr	Pro	Thr	Gly	Arg	Gln	Leu	Thr	Ile
				1205					1210					1215
Phe	Asn	Thr	Gln	Ala	Gln	Ile	Ala	Ile	Gly	Gly	Lys	Asp	Lys	Gly
				1220					1225					1230
Arg	Leu	Phe	Gln	Gly	Gln	Leu	Ser	Gly	Leu	Tyr	Tyr	Asp	Gly	Leu
				1235					1240					1245
Lys	Val	Leu	Asn	Met	Ala	Ala	Glu	Asn	Asn	Pro	Asn	Ile	Lys	Ile
				1250					1255					1260
Asn	Gly	Ser	Val	Arg	Leu	Val	Gly	Glu	Val	Pro	Ser	Ile	Leu	Gly
				1265					1270					1275
Thr	Thr	Gln	Thr	Thr	Ser	Met	Pro	Pro	Glu	Met	Ser	Thr	Thr	Val
				1280					1285					1290
Met	Glu	Thr	Thr	Thr	Thr	Met	Ala	Thr	Thr	Thr	Thr	Arg	Lys	Asn
				1295					1300					1305
Arg	Ser	Thr	Ala	Ser	Ile	Gln	Pro	Thr	Ser	Asp	Asp	Leu	Val	Ser

1310	1315	1320
Ser Ala Glu Cys Ser	Ser Asp Asp Glu Asp	Phe Val Glu Cys Glu
1325	1330	1335
Pro Ser Thr Ala Asn	Pro Thr Glu Pro Gly	Ile Arg Arg Val Pro
1340	1345	1350
Gly Ala Ser Glu Val	Ile Arg Glu Ser Ser	Ser Thr Thr Gly Met
1355	1360	1365
Val Val Gly Ile Val	Ala Ala Ala Ala Leu	Cys Ile Leu Ile Leu
1370	1375	1380
Leu Tyr Ala Met Tyr	Lys Tyr Arg Asn Arg	Asp Glu Gly Ser Tyr
1385	1390	1395
Gln Val Asp Glu Thr	Arg Asn Tyr Ile Ser	Asn Ser Ala Gln Ser
1400	1405	1410
Asn Gly Thr Leu Met	Lys Glu Lys Gln Gln	Ser Ser Lys Ser Gly
1415	1420	1425
His Lys Lys Gln Lys	Asn Lys Asp Arg Glu	Tyr Tyr Val
1430	1435	

<210> 3
 <211> 208
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5398353CD1

<400> 3	
Met Asp Ser Asp Glu Thr Gly Phe Glu His Ser Gly Leu Trp Val	
1 5 10 15	
Ser Val Leu Ala Gly Leu Leu Gly Ala Cys Gln Ala His Pro Ile	
20 25 30	
Pro Asp Ser Ser Pro Leu Leu Gln Phe Gly Gly Gln Val Arg Gln	
35 40 45	
Arg Tyr Leu Tyr Thr Asp Asp Ala Gln Gln Thr Glu Ala His Leu	
50 55 60	
Glu Ile Arg Glu Asp Gly Thr Val Gly Gly Ala Ala Asp Gln Ser	
65 70 75	
Pro Glu Ser Leu Leu Gln Leu Lys Ala Leu Lys Pro Gly Val Ile	
80 85 90	
Gln Ile Leu Gly Val Lys Thr Ser Arg Phe Leu Cys Gln Arg Pro	
95 100 105	
Asp Gly Ala Leu Tyr Gly Ser Leu His Phe Asp Pro Glu Ala Cys	
110 115 120	
Ser Phe Arg Glu Leu Leu Leu Glu Asp Gly Tyr Asn Val Tyr Gln	
125 130 135	
Ser Glu Ala His Gly Leu Pro Leu His Leu Pro Gly Asn Lys Ser	
140 145 150	
Pro His Arg Asp Pro Ala Pro Arg Gly Pro Ala Arg Phe Leu Pro	
155 160 165	
Leu Pro Gly Leu Pro Pro Ala Leu Pro Glu Pro Pro Gly Ile Leu	
170 175 180	
Ala Pro Gln Pro Pro Asp Val Gly Ser Ser Asp Pro Leu Ser Met	
185 190 195	
Val Gly Pro Ser Gln Gly Arg Ser Pro Ser Tyr Ala Ser	
200 205	

<210> 4
 <211> 159
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 71234118CD1

<400> 4

```

Met Ser Phe Phe Asp Asn Ser Thr Ala Ser Ser Asp Asn Trp Lys
 1      5      10      15
Cys Leu Ser Ser Ala Thr Val Cys Thr Leu Phe Leu Phe Ile Ala
 20      25      30
Glu Gln Ser Thr Gly Phe Asn Leu Asp Phe Glu Val Ser Gly Ile
 35      40      45
Tyr Gly Tyr Val Met Leu Asp Gly Met Leu Pro Ser Leu His Ala
 50      55      60
Leu Thr Cys Thr Phe Trp Met Lys Ser Ser Asp Asp Met Asn Tyr
 65      70      75
Gly Thr Pro Ile Ser Tyr Ala Val Asp Asn Gly Ser Asp Asn Thr
 80      85      90
Leu Leu Leu Thr Asp Tyr Asn Gly Trp Val Leu Tyr Val Asn Gly
 95      100      105
Arg Glu Lys Ile Thr Asn Cys Pro Ser Val Asn Asp Gly Arg Trp
 110      115      120
His His Ile Ala Ile Thr Trp Thr Ser Ala Asn Gly Ile Trp Lys
 125      130      135
Val Tyr Ile Asp Gly Lys Leu Ser Asp Gly Gly Ala Gly Leu Ser
 140      145      150
Val Gly Leu Pro Ile Pro Gly Met Phe
 155

```

<210> 5

<211> 500

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 240168CD1

<400> 5

```

Met Ala Ser Val Ala Gln Glu Ser Ala Gly Ser Gln Arg Arg Leu
 1      5      10      15
Pro Pro Arg His Gly Ala Leu Arg Gly Leu Leu Leu Cys Leu
 20      25      30
Trp Leu Pro Ser Gly Arg Ala Ala Leu Pro Pro Ala Ala Pro Leu
 35      40      45
Ser Glu Leu His Ala Gln Leu Ser Gly Val Glu Gln Leu Leu Glu
 50      55      60
Glu Phe Arg Arg Gln Leu Gln Gln Glu Arg Pro Gln Glu Glu Leu
 65      70      75
Glu Leu Glu Leu Arg Ala Gly Gly Gly Pro Gln Glu Asp Cys Pro
 80      85      90
Gly Pro Gly Ser Gly Gly Tyr Ser Ala Met Pro Asp Ala Ile Ile
 95      100      105
Arg Thr Lys Asp Ser Leu Ala Ala Gly Ala Ser Phe Leu Arg Ala
 110      115      120
Pro Ala Ala Val Arg Gly Trp Arg Gln Cys Val Ala Ala Cys Cys
 125      130      135
Ser Glu Pro Arg Cys Ser Val Ala Val Val Glu Leu Pro Arg Arg
 140      145      150
Pro Ala Pro Pro Ala Ala Val Leu Gly Cys Tyr Leu Phe Asn Cys
 155      160      165
Thr Ala Arg Gly Arg Asn Val Cys Lys Phe Ala Leu His Ser Gly
 170      175      180
Tyr Ser Ser Tyr Ser Leu Ser Arg Ala Pro Asp Gly Ala Ala Leu
 185      190      195
Ala Thr Ala Arg Ala Ser Pro Arg Gln Glu Lys Asp Ala Pro Pro
 200      205      210
Leu Ser Lys Ala Gly Gln Asp Val Val Leu His Leu Pro Thr Asp
 215      220      225
Gly Val Val Leu Asp Gly Arg Glu Ser Thr Asp Asp His Ala Ile
 230      235      240
Val Gln Tyr Glu Trp Ala Leu Leu Gln Gly Asp Pro Ser Val Asp
 245      250      255

```



```

Met Lys Val Pro Gln Ser Gly Thr Leu Lys Leu Ser His Leu Gln
260 265
Glu Gly Thr Tyr Thr Phe Gln Leu Thr Val Thr Asp Thr Ala Gly
275 280 285
Gln Arg Ser Ser Asp Asn Val Ser Val Thr Val Leu Arg Ala Ala
290 295 300
Tyr Ser Thr Gly Gly Cys Leu His Thr Cys Ser Arg Tyr His Phe
305 310 315
Phe Cys Asp Asp Gly Cys Cys Ile Asp Ile Thr Leu Ala Cys Asp
320 325 330
Gly Val Gln Gln Cys Pro Asp Gly Ser Asp Glu Asp Phe Cys Gln
335 340 345
Asn Leu Gly Leu Asp Arg Lys Met Val Thr His Thr Ala Ala Ser
350 355 360
Pro Ala Leu Pro Arg Thr Thr Gly Pro Ser Glu Asp Ala Gly Gly
365 370 375
Asp Ser Leu Val Glu Lys Ser Gln Lys Ala Thr Ala Pro Asn Lys
380 385 390
Pro Pro Ala Leu Ser Asn Thr Glu Lys Arg Asn His Ser Ala Phe
395 400 405
Trp Gly Pro Glu Ser Gln Ile Ile Pro Val Met Pro Asp Ser Ser
410 415 420
Ser Ser Gly Lys Asn Arg Lys Glu Glu Ser Tyr Ile Phe Glu Ser
425 430 435
Lys Gly Asp Gly Gly Gly Gly Glu His Pro Ala Pro Glu Thr Gly
440 445 450
Ala Val Leu Pro Leu Ala Leu Gly Leu Ala Ile Thr Ala Leu Leu
455 460 465
Leu Leu Met Val Ala Cys Arg Leu Arg Leu Val Lys Gln Lys Leu
470 475 480
Lys Lys Ala Arg Pro Ile Thr Ser Glu Glu Ser Asp Tyr Leu Ile
485 490 495
Asn Gly Met Tyr Leu
500

```

<210> 6

<211> 455

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7481107CD1

<400> 6

```

Met Asp Thr Pro Arg Val Leu Leu Ser Ala Val Phe Leu Ile Ser
1 5 10 15
Phe Leu Trp Asp Leu Pro Gly Phe Gln Gln Ala Ser Ile Ser Ser
20 25 30
Ser Ser Ser Ser Ala Glu Leu Gly Ser Thr Lys Gly Met Arg Ser
35 40 45
Arg Lys Glu Gly Lys Met Gln Arg Ala Pro Arg Asp Ser Asp Ala
50 55 60
Gly Arg Glu Gly Gln Glu Pro Gln Pro Arg Pro Gln Asp Glu Pro
65 70 75
Arg Ala Gln Gln Pro Arg Ala Gln Glu Pro Pro Gly Arg Gly Pro
80 85 90
Arg Val Val Pro His Glu Tyr Met Leu Ser Ile Tyr Arg Thr Tyr
95 100 105
Ser Ile Ala Glu Lys Leu Gly Ile Asn Ala Ser Phe Phe Gln Ser
110 115 120
Ser Lys Ser Ala Asn Thr Ile Thr Ser Phe Val Asp Arg Gly Leu
125 130 135
Asp Asp Leu Ser His Thr Pro Leu Arg Arg Gln Lys Tyr Leu Phe
140 145 150
Asp Val Ser Met Leu Ser Asp Lys Glu Glu Leu Val Gly Ala Glu
155 160 165

```

```

Leu Arg Leu Phe Arg Gln Ala Pro Ser Ala Pro Trp Gly Pro Pro
170 175 180
Ala Gly Pro Leu His Val Gln Leu Phe Pro Cys Leu Ser Pro Leu
185 190 195
Leu Leu Asp Ala Arg Thr Leu Asp Pro Gln Gly Ala Pro Pro Ala
200 205 210
Gly Trp Glu Val Phe Asp Val Trp Gln Gly Leu Arg His Gln Pro
215 220 225
Trp Lys Gln Leu Cys Leu Glu Leu Arg Ala Ala Trp Gly Glu Leu
230 235 240
Asp Ala Gly Glu Ala Glu Ala Arg Ala Arg Gly Pro Gln Gln Pro
245 250 255
Pro Pro Pro Asp Leu Arg Ser Leu Gly Phe Gly Arg Arg Val Arg
260 265 270
Pro Pro Gln Glu Arg Ala Leu Leu Val Val Phe Thr Arg Ser Gln
275 280 285
Arg Lys Asn Leu Phe Ala Glu Met Arg Glu Gln Leu Gly Ser Ala
290 295 300
Glu Ala Ala Gly Pro Gly Ala Gly Ala Glu Gly Ser Trp Pro Pro
305 310 315
Pro Ser Gly Ala Pro Asp Ala Arg Pro Trp Leu Pro Ser Pro Gly
320 325 330
Arg Arg Arg Arg Arg Thr Ala Phe Ala Ser Arg His Gly Lys Arg
335 340 345
His Gly Lys Lys Ser Arg Leu Arg Cys Ser Lys Lys Pro Leu His
350 355 360
Val Asn Phe Lys Glu Leu Gly Trp Asp Asp Trp Ile Ile Ala Pro
365 370 375
Leu Glu Tyr Glu Ala Tyr His Cys Glu Gly Val Cys Asp Phe Pro
380 385 390
Leu Arg Ser His Leu Glu Pro Thr Asn His Ala Ile Ile Gln Thr
395 400 405
Leu Met Asn Ser Met Asp Pro Gly Ser Thr Pro Pro Ser Cys Cys
410 415 420
Val Pro Thr Lys Leu Thr Pro Ile Ser Ile Leu Tyr Ile Asp Ala
425 430 435
Gly Asn Asn Val Val Tyr Lys Gln Tyr Glu Asp Met Val Val Glu
440 445 450
Ser Cys Gly Cys Arg
455

```

<210> 7

<211> 121

<212> PRT

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7476245CD1

<400> 7

```

Met Ala Arg Arg Ala Gly Gly Ala Arg Met Phe Gly Ser Leu Leu
1 5 10 15
Leu Phe Ala Leu Leu Ala Ala Gly Val Ala Pro Leu Ser Trp Asp
20 25 30
Leu Pro Glu Pro Arg Ser Arg Ala Ser Lys Ile Arg Val His Ser
35 40 45
Arg Gly Asn Leu Trp Ala Thr Gly His Phe Met Gly Lys Lys Ser
50 55 60
Leu Glu Pro Ser Ser Pro Ser Pro Leu Gly Thr Ala Pro His Thr
65 70 75
Ser Leu Arg Asp Gln Arg Leu Gln Leu Ser His Asp Leu Leu Gly
80 85 90
Ile Leu Leu Leu Lys Ala Leu Gly Val Ser Leu Ser Arg Pro
95 100 105
Ala Pro Gln Ile Gln Tyr Arg Arg Leu Leu Val Gln Ile Leu Gln
110 115 120

```

Lys

<210> 8
 <211> 55
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5819744CD1

<400> 8
 Met Thr Ala Ser Glu Cys Gly Thr Gly Asp Gln Ser Cys Arg Arg
 1 5 10 15
 Val Ser Trp Val Val Leu Glu Glu Asp Leu Glu Asp Asp Ala Glu
 20 25 30
 Lys Asp Lys Leu Asn Arg Arg Leu Val Val Leu Leu Gln Leu Leu
 35 40 45
 Tyr Arg Gly Val Lys Pro Ser Ser Ser Val
 50 55

<210> 9
 <211> 545
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5378618CD1

<400> 9
 Met Ala Leu Arg Arg Gly Gly Cys Gly Ala Leu Gly Leu Leu Leu
 1 5 10 15
 Leu Leu Leu Gly Ala Ala Cys Leu Ile Pro Arg Ser Ala Gln Val
 20 25 30
 Arg Arg Leu Ala Arg Cys Pro Ala Thr Cys Ser Cys Thr Lys Glu
 35 40 45
 Ser Ile Ile Cys Val Gly Ser Ser Trp Val Pro Arg Ile Val Pro
 50 55 60
 Gly Asp Ile Ser Ser Leu Ser Leu Val Asn Gly Thr Phe Ser Glu
 65 70 75
 Ile Lys Asp Arg Met Phe Ser His Leu Pro Ser Leu Gln Leu Leu
 80 85 90
 Leu Leu Asn Ser Asn Ser Phe Thr Ile Ile Arg Asp Asp Ala Phe
 95 100 105
 Ala Gly Leu Phe His Leu Glu Tyr Leu Phe Ile Glu Gly Asn Lys
 110 115 120
 Ile Glu Thr Ile Ser Arg Asn Ala Phe Arg Gly Leu Arg Asp Leu
 125 130 135
 Thr His Leu Ser Leu Ala Asn Asn His Ile Lys Ala Leu Pro Arg
 140 145 150
 Asp Val Phe Ser Asp Leu Asp Ser Leu Ile Glu Leu Asp Leu Arg
 155 160 165
 Gly Asn Lys Phe Glu Cys Asp Cys Lys Ala Lys Trp Leu Tyr Leu
 170 175 180
 Trp Leu Lys Met Thr Asn Ser Thr Val Ser Asp Val Leu Cys Ile
 185 190 195
 Gly Pro Pro Glu Tyr Gln Glu Lys Lys Leu Asn Asp Val Thr Ser
 200 205 210
 Phe Asp Tyr Glu Cys Thr Thr Thr Asp Phe Val Val His Gln Thr
 215 220 225
 Leu Pro Tyr Gln Ser Val Ser Val Asp Thr Phe Asn Ser Lys Asn
 230 235 240
 Asp Val Tyr Val Ala Ile Ala Gln Pro Ser Met Glu Asn Cys Met
 245 250 255
 Val Leu Glu Trp Asp His Ile Glu Met Asn Phe Arg Ser Tyr Asp

Asn Ile Thr Gly	260	Gln Ser Ile Val Gly	265	Cys Lys Ala Ile Leu	270
Asp Asp Gln Val	275	Phe Val Val Val Ala	280	Gln Leu Phe Gly Gly	285
His Ile Tyr Lys	290	Tyr Asp Glu Ser Trp	295	Thr Lys Phe Val Lys	300
Gln Asp Ile Glu	305	Val Ser Arg Ile Ser	310	Lys Pro Asn Asp Ile	315
Leu Phe Gln Ile	320	Asp Asp Glu Thr Phe	325	Phe Val Ile Ala Asp	330
Ser Lys Ala Gly	335	Leu Ser Thr Val Tyr	340	Lys Trp Asn Ser Lys	345
Phe Tyr Ser Tyr	350	Gln Ser Leu His Glu	355	Trp Phe Arg Asp Thr	360
Ala Glu Phe Val	365	Asp Ile Asp Gly Lys	370	Ser His Leu Ile Leu	375
Ser Arg Ser Gln	380	Val Pro Ile Ile Leu	385	Gln Trp Asn Lys Ser	390
Lys Lys Phe Val	395	Pro His Gly Asp Ile	400	Pro Asn Met Glu Asp	405
Leu Ala Val Lys	410	Ser Phe Arg Met Gln	415	Asn Thr Leu Tyr Leu	420
Leu Thr Arg Phe	425	Ile Gly Asp Ser Arg	430	Val Met Arg Trp Asn	435
Lys Gln Phe Val	440	Glu Ile Gln Ala Leu	445	Pro Ser Arg Gly Ala	450
Thr Leu Gln Pro	455	Phe Ser Phe Lys Asp	460	Asn His Tyr Leu Ala	465
Gly Ser Asp Tyr	470	Thr Phe Ser Gln Ile	475	Tyr Gln Trp Asp Lys	480
Lys Gln Leu Phe	485	Lys Lys Phe Lys Glu	490	Ile Tyr Val Gln Ala	495
Arg Ser Phe Thr	500	Ala Val Ser Thr Asp	505	Arg Arg Asp Phe Phe	510
Ala Ser Ser Phe	515	Lys Gly Lys Thr Lys	520	Ile Phe Glu His Ile	525
Val Asp Leu Ser	530	Leu	535		540
	545				

<210> 10
 <211> 1374
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 1657368CB1

<400> 10
 ccaagttcgt gggctctctc agaagtcctc aggacggagc agaggtggcc ggccgggccg 60
 gctgactgcg cctctgcttt ctttcataa ctttttcttt cggactcgaa tcacggctgc 120
 tgccaagggg ctagtccggg acactagggt gcccgaacgc gctgatgccc cgagtgcctg 180
 cagggcttcc cgtaaccat gctgccggcg ccggcgcccg cagctgcctt ggcgtgcctc 240
 gtgctcctgc tactgctggt ggtgctgacg ccgccccga ccggcgcaag gccatcccca 300
 ggcccagatt acctgcggcg cggctggatg cggctgctag cggagggcga gggctgcgct 360
 ccctgccggc cagaagagtg cgccgcggcg cggggctgcc tggcgggcag ggtgcgcgac 420
 gcgtgcggct gctgctggga atgcgccaac ctcgagggcc agctctgcga cctggacccc 480
 agtgctcact tctacgggca ctgcggcgag cagcttgagt gccggctgga cacaggcggc 540
 gacctgagcc gcggagaggt gccggaacct ctgtgtgcct gtcgttcgca gagtccgctc 600
 tgcgggtccg acggtcacac ctactcccag atctgccgcc tgcaggaggc ggcccgcgct 660
 cggcccgatg ccaacctcac tgtggcacac ccggggccct gcgaatcggg gccccagatc 720
 gtgtcacatc catatgacac ttggaatgtg acagggcagg atgtgatctt tggctgtgaa 780
 gtgtttgcct accccatggc ctccatcgag tggaggaagg atggcttgga catccagctg 840
 ccaggggatg accccacat ctctgtgcag tttagggggtg gaccccagag gtttgagggtg 900
 actggctggc tgcagatcca ggctgtgcgt cccagtgatg agggcactta ccgctgcctt 960
 ggccgcaatg ccctgggtca agtggaggcc cctgctagct tgacagtgc caccctgac 1020

cagctgaact	ctacaggcat	ccccagctg	cgatcactaa	acctgggtcc	tgaggaggag	1080
gctgagagt	aagagaatga	cgattactac	taggtccaga	gctctggccc	atgggggtgg	1140
gtgagcggc	atagtgttca	tccctgctct	tgaaaagacc	tgaaaagggg	agcagggtcc	1200
cttcatcgac	tgctttcatg	ctgtcagtag	ggatgatcat	gggaggccta	tttgactcca	1260
aggtagcagt	gtggtaggat	agagacaaaa	gctggaggaa	ggtagggaga	gaaactgaga	1320
ccaggaccgc	tgggggtacaa	agggggccat	gcaggagata	gcctggccag	tagg	1374

<210> 11

<211> 4541

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 4028972CB1

<400> 11

tctgtgtgtg	tgtgtgccttc	ctcctgtgtg	ctttctgttc	ccccatctct	gtcttgttct	60
tcccacttct	attgccaaag	ggagagatcc	tctccgggct	gttccctggc	ctgtctgtct	120
ctccgggctc	tgtcccagca	gcgacaatga	gtcccacact	ccactcgggt	ttcttcaccc	180
tgaaggtcag	catcctgctg	gggtccctgc	tggggctctg	cctgggcctt	gagttcatgg	240
gcctcccca	ccagtgggac	cgctacctcc	gctgggatgc	cagcacacgc	agtacactga	300
gtttccagtt	caagaccaac	gtctctacgg	ggctgtctct	ctacctggat	gatggcggcg	360
tctgcgactt	ctatgccttc	tccctgggtg	atggccgcgt	tcagctccgc	ttcagctccg	420
actgtgccga	gactgccgtg	ctgtccaaca	agcagggtga	tgacagcagc	tggcacttcc	480
tcattggtgag	ccgtgacccg	ctgcgcacgg	tgtctgatgt	tgatggcgag	ggccagtctg	540
gggagctgca	gccccagcgg	ccctacatgg	atgtggctcag	tgacttgttc	cttgggtggag	600
tcctactactga	catacgacct	tctgccctga	cccttgatgg	agttcaggcc	atgcccggct	660
tcaagggggt	aattctggat	ctcaagtatg	gaaactcgga	gcctcggcct	ctggggagcc	720
gggggtgtcca	gatggatgcc	gagggaccct	gtgggtgagc	tcctctgtgaa	aatggtggga	780
tctgctttct	ctggacggc	caccccacct	gtgactgttc	taccactggc	tatggtggca	840
agctctgtct	agaagatgtc	agtcaagatc	caggcctctc	ccacctcatg	atgagtgaac	900
aaggtagaag	taaagctcga	gaggagaatg	tggccacttt	ccgaggctca	gagtatctgt	960
gctacgacct	gtctcagaac	ccgatccaga	gcagcagtga	tgaatcacc	ctctccttta	1020
agacctggca	gcgtaacggc	ctcatctgc	acacgggcaa	gtcggctgac	tatgtcaacc	1080
tggctctgaa	ggatggtgag	gtctccttgg	tcattaacct	ggggctcggg	gcctttgagg	1140
ccattgttga	gccagtgaat	ggaaaattca	acgacaacgc	ctggcatgat	gtcaaatgta	1200
cacgcaacct	ccggcagggt	acaatctctg	tggatggcat	tcttaccacg	acgggctaca	1260
ctcaagagga	ctataccatg	ctgggctcgg	acgaacttct	ctatgtagga	ggaagcccaa	1320
gtaccgctga	cttgccctggc	tcccctgtca	gcaacaactt	catgggctgc	cttaaaggagg	1380
ttgtttataa	gaataatgac	atccgtcttg	agctgtctcg	cctggcccg	attgcccaga	1440
ccaagtgtaa	aatctatggc	gaagtgtgtg	ttaagtgtga	gaatgtggcc	acactggacc	1500
ccatcaactt	tgagacccca	gaggcttaca	tcagcttgcc	caagtggaac	actaaacgta	1560
tgggctccat	ctcctttgac	ttccgcacca	cagagcccaa	tggcctgata	ctcttcaact	1620
atggaaagcc	ccaagagagg	aaggatgtct	ggagccagaa	gaatacaaaa	gtagacttct	1680
ttgcctggga	actcctcgat	ggcaacctgt	acttgcctgt	tgacatgggc	tctggcaca	1740
tcaaagtga	agccactcag	aagaaagcca	atgatgggga	atggtaccat	gtggacattc	1800
agcgagatgg	cagatcaggt	actatatcag	tgaacagcag	gcgcacgcca	ttcaccccca	1860
gtggggagag	cgagatcctg	gacctggaag	gagacatgta	cctgggagg	ctgccggaga	1920
accgtgctgg	ccttattctc	cccaccgagc	tgtggactgc	catgctcaac	tatggctacg	1980
tgggctgcat	ccgcgacct	ttcattgatg	ggcgcagcaa	gaacattcga	cagctggcag	2040
agatgcagaa	tgctgccccg	gtcaagtcct	cctgttcacg	gatgagtggc	aagcagtgtg	2100
acagctaccc	ctgcaagaat	aatgctgtgt	gcaaggacgg	ctggaaccgc	ttcatctgcg	2160
actgcaccgg	caccggatac	tggggaagaa	cctgcgaaag	ggaggcatcc	atcctgagct	2220
atgatggtag	catgtacatg	aagatcatca	tgcccatggg	catgcatact	gaggcagagg	2280
atgtgtcctt	ccgcttcgat	tcccagcgag	cttatgggct	gctggtggct	acgacctcca	2340
gggactctgc	cgacaccctg	cgtctggagc	tggatggggg	gcgtgtcaag	ctcatggtta	2400
acttagactg	tatcaggata	aactgttaact	ccagcaaaag	accagagacc	ttgtatgcag	2460
ggcagaagct	caatgacaac	gagtggcaca	ccgttcgggt	ggtgcccaga	ggaaaaagcc	2520
taaagttaac	cgtggatgat	gatgtggctg	agggtacaat	ggtgggagac	catacccggt	2580
tggagtcca	caacattgaa	acgggaatca	tgactgagaa	acgctacatc	tccgttgtcc	2640
cctccagctt	tattggccat	ctgcagagcc	tcatgtttaa	tggccttctc	tacattgact	2700
tgtgcaaaaa	tggtgacatt	gattattgtg	agctgaaggg	tcgttttggg	ctgaggaaca	2760
tcactgcctga	ccctgtcacc	tttaagacca	agagcagcta	cctgagcctt	gccactcttc	2820
aggcttacac	ctccatgcac	ctcttcttcc	agttcaagac	cacctcacca	gatggcttca	2880
ttctcttcaa	tagtggtgat	ggcaatgact	tcattgcagt	cgagcttgct	aaggggtata	2940
tacactacgt	ttttgacctc	ggaaacggtc	ccaatgtgat	caaaggcaac	agtgaccgcc	3000

```

cctgaatga caaccagtgg cacaatgtcg tcatactcg ggacaatagt aacactcata 3060
gcctgaaagt ggacacccaaa gtggtcactc aggttatcaa tgggtgccaaa aatctggatt 3120
tgaaaggtga tctctatatg gctgggtctgg cccaaggcat gtacagcaac ctcccaaagc 3180
tcgtggcctc tcgagatggc ttccagggct gtctagcatc agtggacttg aatggacgcc 3240
tgccagacct catcaatgat gctcttcacg ggagcggaca gatcgagcgt ggctgtgaag 3300
gaccaggtac cacttgccag gaagattcat gtgccaaacca gggggtctgc atgcaacaat 3360
gggagggtct cactgtgat tgtctatga cctcttatte tggaaaccag tgcaatgatc 3420
ctggcgctac gtacatcttt gggaaaagtg gtgggcttat cctctacacc tggccagcca 3480
atgacaggcc cagcacgcgg tctgaccgcc ttgccgtggg cttcagcacc actgtgaagg 3540
atggcatctt ggtccgcacg gacagtgtct caggacttgg tgacttcctc cagcttcaca 3600
tagaacaggg gaaaattgga gttgtcttca acattggcac agttgacatc tccatcaaag 3660
aggagagaac cctgttaaag gacggcaaat accatgtggt acgcttcacc aggaacggcg 3720
gcaacgccac cctgcagggt gacaactggc cagtgaatga acattatcct acaggccggc 3780
agttaacctt cttcaacact caggcgcaaa tagccattgg tggaaaggac aaaggacgcc 3840
tcttccaagg ccaactctct gggctctatt atgatggttt gaaagtactg aacatggcgg 3900
ctgagaacaa ccccaatatt aaaatcaatg gaagtgttct gctgggttgg gaagtcccat 3960
caattttggg aacaacacag acgacctcca tgccaccaga aatgtctact actgtcatgg 4020
aaaccactac tacaatggcg actaccacaa cccgtaagaa tcgctctaca gccagcattc 4080
agccaacatc agatgatctt gtttcatctg ctgaatgttc aagtgtatga gaagactttg 4140
ttgaatgtga gccgagtaca gcaaacccca cggagccggg aatcagacgg gttccggggg 4200
cctcagaggt gatccggggg tcgagcagca caacagggat ggtcgtcggc attgtggctg 4260
ctgccgccct ctgcatcttg atcctcctgt acgccatgta caagtacagg aacagggacg 4320
aggggtccta tcaagtggac gagacgcgga actacatcag caactccgcc cagagcaacg 4380
gcacgctcat gaaggagaag cagcagagct cgaagagcgg ccacaagaaa cagaaaaaca 4440
aggcagggg gtattacgtg taaacatgcg aacactgtct acacgcgagt tttcacagtt 4500
atctctatcc acgcctatga atctttggac ggtgagatct c 4541

```

<210> 12
 <211> 1117
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5398353CB1

```

<400> 12
ctctgcctag cacatcccc ctcccacctc ctcatecaca aaatgtatag gtttgcataa 60
aataaggtgg aaaattagac agcagcgaga tcatgaaggg tgttgaatga caccgagtaa 120
atacacttta acctatagaa ttttaaggca aaaagtgagc tatgacgtct gcaagcaagc 180
ggtaagtaaa gtccggaatc cgggttcgag gctgtcagct gaggatccag ccgaaagagg 240
agccaggcac tcaggccacc tgagtctact cacctggaca actggaatct ggcaccaatt 300
ctaaaccact cagcttctcc gagctcacac cccggagatc acctgaggac ccgagccatt 360
gatggactcg gacgagaccg ggttcgagca ctcaggactg tgggtttctg tgctggctgg 420
tctgctggga gcctgccagg cacaccccat ccctgactcc agtctctctc tgcaattcgg 480
gggccaagtc cggcagcggg acctctacac agatgatgcc cagcagacag aagcccacct 540
ggagatcagg gaggatggga cgggtggggg cgtgctgac cagagccccg aaagtctcct 600
gcagctgaaa gccttgaagc cgggagttat tcaaattctg ggagtcaaga catccagggt 660
cctgtgccag cggccagatg gggcctgtga tggatcgctc cactttgacc ctgaggcctg 720
cagcttccgg gagctgcttc ttgaggacgg atacaatgtt taccagtcgg aagcccacgg 780
cctccgctg cacttgccag ggaacaagtc cccacaccgg gacctgcac cccgaggacc 840
agctcgtctt ctgccactac caggcctgcc ccccgcactc ccggagccac ccggaatcct 900
ggccccccag ccccccgatg tgggctctct ggaccctctg agcatgggtg gaccttccca 960
gggccgaagc cccagctacg cttcctgaag ccagagcgtg tttactatga catctcctct 1020
ttattttatta ggttattttat cttattttat tttttttttt cttacttgag ataataaaga 1080
gttcacagag agaaaaaaaa aaaaaaaaaa aaaaaaa 1117

```

<210> 13
 <211> 2460
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 71234118CB1

<220>

<221> unsure

<222> 2443 :

<223> a, t, c, g, or other

<400> 13

```

cctgtgtgca gctggcttca caggatcaca ctgtgaattg aacatcaatg aatgtcagtc 60
taatccatgt agaaatcagg ccacctgtgt ggatgaatta aattcataca gttgtaaatg 120
tcagccagga ttttcaggca aaagggtgtga aacagggtatg tatcaactca gtgttattaa 180
taacaatact aacaatagta atataataac aataatacca attttgctg caccaagcag 240
tgccgtagg ccaaaaacga tgccaagcac tttgtactca ttaatcttc acagccatcc 300
aacaactag gtactactac tattatcttc agtggcacag atgatgaaat taaagcttaa 360
aaattaggg gggaatatga gaaaattgaa tttagtagaa ctcttcatgt tctacttatt 420
tataatagaa atgatacttt gataagccgc ttgacttcag agtgagact ccatggacat 480
gtttgagaat cagcagatca aatctaaata ctttctcagg agactggagt cttttagctg 540
taagacttgt ttatttgagt ggtctgtgca ggtttggggc agactggaat tgctcatgaa 600
atagggaaaa gccagagcct tgagtagggc agcatgttgc cettgcagca cctcctcctt 660
ctctcctgtg acatcaatta cttcctctct actgagtcac tctcatcagc atacggacat 720
gcttttagtat ctcttggtt tagtagcatt tcaatttcac ctctccttcc agctacttcc 780
tgtttctctg ctctcttcca ctattgaatg ttttctctat cttatatact cactggttat 840
tttattttaa agaatgcact atgtattttg taaataatat gtatttaatt ttgaaactgc 900
ttcttagata atcatattca ctgaaatgag cttctttgac aatagcactg cttcctctga 960
caattggaaa tgtctttcaa gtgccacagt ctgcacactt ttcttattoa ttgcagaaca 1020
gtctacaggc tttaacctgg attttgaagt ttctggcatc tatggatatg tcatgctaga 1080
tgccatgctc ccactctccc atgctctaac ctgtaccttc tggatgaaat cctctgacga 1140
catgaactat ggaacaccaa tctcctatgc agttgataac ggcagcgaca ataccttgct 1200
cctgactgat tataacggct gggttcttta tgtgaatggc agggaaaaga taacaaactg 1260
tccctcgggtg aatgatggca gatggcatca tattgcaatc acttgacaa gtgccaatgg 1320
catctggaaa ctctatatcg atgggaaatt atctgacggg ggtgctggcc tctctgttgg 1380
tttgcccata cctggtatgt tttacaaaa tgagtataat tctgttgga tctagagggtg 1440
atgctgctta attgacatc aggttttgta gcattgttgt tcttggttcc aaacaaagat 1500
taacagctat ctctctcatt tctactccc tgattatcct aactgtgggt tattagaagc 1560
tttttctcta agtaagctgt ttgcctctct gtgtgagaaa tgcttttgaa gatctacatt 1620
aagactacca agaagcatgc tgaactagca gctgggttgc aaactgtctt ctaaggagat 1680
taattactgg tattttttgt ttattttaag gattgttttg ttttgtaatc agtatataag 1740
caggaccact gctatctatc ttttatggcc cttttcaaat agatcttggc ttgtcttcta 1800
aaaaatactt tttaataaat atcccaggag ttttagtttg aagaattgca aactaaattt 1860
cattaggtga ccagggttct cagcattgta catggcaata agcttgttct atctggatac 1920
atctatagaa gtatatgcat atgacattat tttttacttt gctcatttgt ttcaggata 1980
ctaaaagaaa attaaataga caaaaatgaa ttaaaagtatt agaaactgac taatacattc 2040
tagaagtaag ttgtatgaaa ctaggatacg tataactaata aatatagcac aaaaatacta 2100
aacttcataa aaatgcgggtg agatgacctc tgccaagcct caaatacaat acttatttta 2160
tttattagaa attattggcc aggtacgggt gctcacgccc gtaatcccag cactttggaa 2220
ggctgaggag ggtggatcac gaggtcagga gttggagacc gccctggaca atatggtgaa 2280
gccctatctc tactaaaaac gcacagaaaa attagccagg catggtggta gatgcctgta 2340
attccaggca ctcaggaggc tgaggcagga gaatcgcttg aaccaggag gcgaagattg 2400
ccgtgagcgg agggcatgcg actcgaggcc aagaatgggg agnggcagg agagggacgg 2460

```

<210> 14

<211> 2601

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 240168CB1

<400> 14

```

gggttcgggg gcgcgcgct gtgaggccgg ggcttagagc cagccgcggc cgccgaggag 60
gggcccaggc ccgcgctcg cccgcgtccc cgccttcttc ccgcgctcag cccgcctcg 120
gctcgctgcc cttggctctc gtcgccatgg cctccgtcgc ccaggagagc ggggctcgc 180
agcgcgggct accgcgcgt cacggggcgc tgcgggggct gctactgctc tgctgtggc 240
tgccaagcgg ccgtgcggcc ttgccgcccg cggcgccgct gtccgaactg cagcgcagc 300
tgtggggcgt ggagcagctg ctggaggagt tccgcggca actgcagcag gagcggcctc 360
aggagagct ctgcgcgccc gggcgggccc ccaggaggac tgccggggcc 420
cgggcagcgg cggctacagc gcaatgcctg acgccatcat ccgcaccaag gactccctgg 480
cggcgggtgc cagcttctct cgggcgcggc cggccgtgag gggctggcgg caatgcgtgg 540
cggcctgctg ctccgagccc cgctgctccc tggcgtgggt ggagctgccc cggcgccccg 600

```

```

cgcccccggc agccgtgctc ggctgctacc tcttcaactg cagggcggcg gggcgcaacg 660
tctgcaagtt cgcgtgcac agcggctaca gcagctacag cctcagccgc gcgcgggacg 720
gcgccgcctg ggcaccgcg cgcgcctcgc cccggcagga aaaggatgcg cctccactta 780
gcaaggctgg gcaggatgtg gttctgcate tgcccacaga cgggggtggt ctagacggcc 840
gcgagagcac agatgaccac gccatcgctc agtatgagtg ggcactgctg caggggggacc 900
cgtcagtggg catgaagggt cctcaatcag gaacctgaa gctgtccac ctacaggagg 960
gaacctacac ctccagctg accgtgacgg acactgccg gcagagaagc tctgacaacg 1020
tgtcagtgc agtgcttcgc gcagcctact ccacaggagg atgtttgcac acttgctcac 1080
gctaccactt cttctgtgac gatggctgct gcattgacat cagctcgcg tcgatggag 1140
tgcagcagtg tccctgatggg tctgatgaag acttctgcca gaatctgggc ctggaccgca 1200
agatggtaac ccacacggca gctagtcccg cctcgccaag aaccacaggg ccgatgaag 1260
atgcaggggg tgactccttg gtgaaaagt ctcagaaagc cactgcccc aacaagccac 1320
ctgcattatc aaacacagag aagaggaatc attccgcct ttggggacca gagagtcaaa 1380
tcattcctgt gatgccagat agtagttcct caggaagaa cagaaaagag gaaagttata 1440
tattttgagtc aaagggatg ggaggaggag gggaacacc agcccagaa acagggtgcag 1500
tgctaccctt ggcgtgggt ttggctatca ctgctctgct gcttctcatg gttgcatgcc 1560
gactacgact ggtgaaacag aaactgaaaa aagctcgctc cattacatct gaggaatcgg 1620
actacctcat aatgggatg tatctatagt aatgtaatt caataccttg gggcagggac 1680
atgttttgtt tataatttat acatctatta agttctggat atttacagct tcttttgttt 1740
ttaattgggc cagaagattc tgcaaatccc aaatctttct ttattattta ttgtaaaaaa 1800
agtttcttta gaagtcataa aataatttga aatttagaga ggaattcatg attaaagatt 1860
cctaaaaata taattctgat ttatgtaagc tgcctgaa aatagaaatg tgtacttagc 1920
tgagagaaaa ttcagcatct caggaggtgg tattaggatg actgtgttaa cccattacct 1980
tttagaagcc aactgttggc ccttaccat gctggactgc tataggccca gcttccccct 2040
gttctgtggc cctttcttc tgagagtggg tctcaagcaa gttcatgcct tcaatcagat gttacttagg 2160
gtgggtatac cttaaattata aacctatgt acaagtcagt aagccttagg gaaggtgagt 2220
gtgggtcctt cctaaccct ctgacgtcat gtcatatagg tggctgcctc cttagactga 2280
cctttgggag aaaaaaaccc cagactttga attagtaaca gctctaagat ggtcatgcag 2340
tgagatagga aatcaagatg gaagcagaga atctggcatg ccaaaaacta acagaaactt 2400
agttgaaggc aaagagagca aggagaacgt ttaatactc attacatcaa atcaacactg 2460
ctccatgggt aggcacagc aactcattta tatatatata tataggcttt gttgatgaaa 2520
aacaacaatt gaagagagga cgttgagtgg attcctgggt acagcttttg taaaaatgtc 2580
accatggctt tcatccaatg g 2601

```

<210> 15

<211> 2791

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<223> Incyte ID No: 7481107CB1

<400> 15

```

cttggggaag gaggaagtcc tgcaggcggg agggaaagaa gagagggaag atggggatgc 60
agtggaggcg gggggcaggc cgcgagagg agaggatccc gggagcagac gaagaagtgg 120
agcagctaaa gtctgcgtca gaagagggtg gggactgcga gaggagaggc tggggcctgc 180
aggggagcgc agcagctttt agcatcgatc caaactctaa agactcgtgg cctttgcctg 240
acctcgaggg tcgggaatag acgcctgtct ttgtggagag cgatacccaa ccgagaaaat 300
ggggctgttc cgagctgggc cctgcgcctg gccaggggcg aggtctctct ggctccgggc 360
tgccccctga ggggcgcaaa cggcaggcct ggcggttggg gccaggagg gaaggtacca 420
ccgccccgag gcagcacgca gcctgcagca gaggcgctg ctccgagctg tctcttgggg 480
gcgcgcgcgc cgcttccctc ctccggggcc gctcgctccc aggaaagtgg aggggctgg 540
cgaggaccga gagccggggc cgcgtgcgg agggaccaca cctccgggag ttcgaggggg 600
accttggcgc ggcgggccag cctttgctcc cgggccagg gccggcagcg cccgccttcc 660
ccgggtcagc gcttgcggcc cgcgcgcgc gcacgcgcg gcaaccccg gcgctccc 720
cgggggcgct gcgtcttct gccacaccg gcacgcgcg cccctctccc ccacacctcc 780
ggcccgcaac acccggtct cctcccacc tcccccccc tctctgccc tccctcccca 840
ttcttccct cccggcgagg ggcgggagg ggcgtggcg ggccgggggt tgtgtggctg 900
ggacccggct cctgcactc cgagtccgc cgaggagcg ggcccggcc gctgtccagc 960
cgctccgtgc cccgcgcgtc ctgcgcgcc gccaccgct cctggggaga cgcagccact 1020
tgcccgccat ggatactccc agggctctgc tctcgccgt cttctctatc agttttctgt 1080
gggattttcc cggtttccag caggcttcca tctcactct ctcgtctgct gccgagctgg 1140
gttccaccaa ggcgatgcga agccgcaag atggcaagat gcagcgggcg ccgcgcgaca 1200
gtgacgcggg ccgggagggc caggaaccac agccgcggcc tcaggacgaa ccccgggctc 1260
agcagccccg ggcgcaggag ccgccaggca ggggtccgc cgtggtgccc cagagtaca 1320

```



```

tgctgtcaat ctacaggact tactccatcg ctgagaagct gggcatcaat gccagctttt 1380
tccagtcttc caagtccgct aatacagatca ccagctttgt agacagggga ctacagcate 1440
tctcgcacac tectctccgg agacagaagt atttgtttga tgtgtccatg ctctcagaca 1500
aagaagagct ggtgggcgcg gagctcgccg tctttcgcca ggcgccctca gcgccctggg 1560
ggccaccagc cgggcccgtc cacgtgcagc tcttcccttg cctttcgcce ctactgctgg 1620
acgcgcggac cctggaccgg cagggggcgc gcgcggccgg ctgggaagtc ttcgacgtgt 1680
ggcagggcct gcgccaccag ccctgggaag agctgtgctt ggagctgcgg gccgcatggg 1740
gagagctgga gcgcggggag gccgagggcg gcgcgcgggg accccagcaa ccgccgccc 1800
cggacctgcg gagtctgggc ttcggccgga ggggtgcggc tccccaggag cgggcccctg 1860
tggtggtatt caccagatcc cagcgcaaga acctgttcgc agagatgcgc gagcagctgg 1920
gctcggccga ggctgcgggc ccgggcgcgg gcgcgcaggg gtcgtggccg ccgccgtcgg 1980
gcgccccgga tgccaggcct tggctgccct cgccccggcg ccggcggcgg cgcacggcct 2040
tcgccagtcg ccatggcaag cggcacggca agaagtccag gctacgctgc agcaagaagc 2100
ccctgcacgt gaacttcaag gagctgggct gggacgactg gattatcgcg cccctggagt 2160
acgaggccta tcactgcgag ggtgtatgcg acttcccgtc gcgctgcac ctggagccca 2220
ccaaccacgc catcatccag acgctgatga actccatgga ccccggtccc accccgccca 2280
gctgctgctg gccacacaaa ttgactccca tcagcattct atacatcgac gcgggcaata 2340
atgtggctta caagcagtac gaggacatgg tgggtggagt gtgcggtcgc aggtagcggg 2400
gcctttcccg ccgccttggc ccggaaccaaa ggtgggccaag ggtccgcctt gcaggggagg 2460
cctggctgca gagaggcggg ggaggaagct ggcgctgggg gaggtcgagg gtgaggggaa 2520
agcctggatg tgagagccgg tgggagagaa gggagcgag acttcccagt aacttctacc 2580
tgccagccca gagggaaata tggattttca caccttgcct ggacaccctg gaaaaacaag 2640
ccaaggagga tttctttagt tctgtttctc tctctctctc tctctctctc tctctctctc 2700
tctctctctc tctctctctc tatcagtgtg tctgtgtaat cccatgtgtg tcatacagct 2760
cgagatataa tgcgccagac ggcacaacct c

```

<210> 16
 <211> 709
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 7476245CB1

```

<400> 16
ctcagagccgc ggcgccgaac gaagccgcgg cccgggcaca gccatggccc ggccggcggg 60
gggcgctcgg atgtttggca gcctcctgct cttgcgccctg ctcgctgccg gcgtgcgcc 120
gctcagctgg gatctcccgg agccccgcag ccgagccagc aagatccgag tgcaactcgg 180
aggcaacctc tgggccaccg gtcacttcat gggcaagaag agtctggagc cttccagccc 240
atccccattg gggacagctc cccacacctc cctgagggag cagcgactgc agctgagtca 300
tgacttgctc ggaatcctcc tgctaaagaa ggccttgggc gtgagcctca gccgcccg 360
accccaaatc cagtaacagga ggctgctggt acaataatcg cagaaatgac accaataatg 420
gggcagacac aacagcgtgg cttagattgt gccacccag ggaaggtgct gaatgggacc 480
ctggttgatg ccccatctgg atgtaaatcc tgagctcaaa tctctgttac tccattactg 540
tgatttctgg ctgggtcacc agaaatatcg ctgatgcaga cacagattat gttcctgctg 600
tatttctgct ttcctgttg aattggtgaa taaaaccttg ctctttacat aaaaaaaaaa 660
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaacaaaaaa aaaaaaaaaa 709

```

<210> 17
 <211> 753
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5819744CB1

```

<400> 17
ctggacctgg gaccatgtct tgtattgcca gcacagggcc ctgaaacaca agagatattg 60
ggcaagggat agaccagtgg aagagaggta gaaagcaagt ccagagataa agatttagaa 120
aggctgcaga ctaaaatgac tgcttcagaa tgtgggacag gagaccagag ctgcagaaga 180
gtcagctggg tggatttggg ggaggacctg gaagatgacg ctgagaaaga caagcttaac 240
aggagactgg ttgttctctt gcagctgctt tacaggggag tgaagccttc ctcctccgct 300
tgatgctggg cctgccttcc ctgctgtgca gctactctgt gtctgtgggt ctagtacctt 360
catttctcat tatgttcaca caagtcaggt tgttggggag ctgagctctt acagtcccat 420
gggggaagtg gagagccaaa ggtctcactt tttgtgattc tttggagatt ttatgcacc 480

```

tagaaatattg	agtttttccct	gggtgtcttct	gggactggta	ctgcataccc	agggtcacc	540
tggttggggg	cttttccaag	cggagtttgt	gatcataatg	agttatatatt	ttcctctttt	600
ctggataaga	acaatatcct	tgccattcag	tctccaaata	tagcatgtgg	ttatgataga	660
agaatatcct	tatgttagga	aagaaattgg	cagtctcaat	gaggcagaaa	tgtacaattt	720
agaaaacatg	tataaattac	aacagcgggt	gta			753

<210> 18
 <211> 3413
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <223> Incyte ID No: 5378618CB1

<400> 18

cgtctgcagc	ggcgccgggt	ccgagcgcgc	ggcgcgccgc	tgggggtcgg	ggcccgggcg	60
gggagcgggg	accgggcatg	gcgctgcgga	gaggcgccgc	cggagcgctc	gggctgtctg	120
tgctgtctgt	gggcgcgcgc	tgccctgatac	cgcggagcgc	gcagggtgagg	cggctggcgc	180
gctgccccgc	cacttgcagc	tgtaccaagg	agtctatcat	ctgcgtgggc	tcttctctgg	240
tgcccaggat	cgtgcggggc	gacatcagct	ccctgagcct	ggtaaatggg	acgttttcag	300
aaatcaagg	ccgaatgttt	tcccatctgc	cttctctgca	gctgctattg	ctgaattcta	360
actcattcac	gcatcatcgg	gatgatgctt	ttgctggact	ttttcatctt	gaataacctg	420
tcattgaagg	gaacaaaata	gaaaccattt	caagaaatgc	ctttcgtggc	ctccgtgacc	480
tgactcacct	ttctttggcc	aataaccaca	taaaagcact	accaagggat	gtcttcagtg	540
atthagactc	tctgattgaa	ctagatttga	ggggtaataa	atttgaatgt	gactgcaaag	600
ccaagtggct	atacctgtgg	ttgaagatga	caaattccac	cgtttctgat	gtgctgtgta	660
ttggtccacc	agagtatcag	gaaaagaagc	taaatgacgt	gaccagcttt	gactatgaat	720
gcacaactac	agattttgtt	gttcatcaga	ctttacccta	ccagtcgggt	tcagtgagta	780
cgttcaactc	caagaacgat	gtgtacgtgg	ccatcgcgca	gcccagcatg	gagaactgca	840
tggtgtctgga	gtgggaccac	attgaaatga	atltccggag	ctatgacaac	attacagggtc	900
agtcctatcg	gggctgtgaa	gccattctca	tcgatgatca	ggtctttgtg	gtggtagccc	960
agctcttcgg	tggtctctac	atltacaaat	acgacgagag	ttggaccaaa	tttgcataat	1020
tccaagacat	agaggtctct	cgcatttcca	agcccaatga	catcgagctg	tttcagatcg	1080
acgacgagac	gttctttgtc	atcgacagca	gctcaaaggc	tggtctgtcc	acagtttata	1140
aatggaacag	caaaggattc	tattcttacc	agtcactgca	cagatgggtc	agggacacgg	1200
atgcggagtt	tggtgatata	gatggaaaat	cgcatctcat	cctgtccagc	cgctcccagg	1260
tcccatcat	cctccagtgg	aataaaagct	ctaagaagtt	tgtcccccac	ggtgacatcc	1320
ccaacatgga	ggacgtactg	gctgtgaaga	gcttccgaat	gcaaaaatac	ctctaccttt	1380
cccttaccgc	cttcacggg	gactccggg	tcattgagtg	gaacagtaag	cagtttctgg	1440
agatccaagc	tcttccatcc	cggggggcca	tgaccctgca	gcccttttct	tttaagata	1500
atcactacct	ggccctgggg	agtgaactata	cattctctca	gatataccag	tgggataaag	1560
agaagcagct	attcaaaaag	tttaaggaga	tttacgtgca	ggcgccctcg	tcattcacag	1620
ctgtctccac	cgacaggaga	gatttctttt	ttgcatccag	tttcaaaggg	aaaacaaaga	1680
tttttgaaca	tataattgtt	gacttaagtt	tgtgaagggt	tggtgggtga	aactaagaga	1740
aatgtagcat	tagctctcac	aaaagaggac	caagaaaaat	caacaaacaa	atcaaagcca	1800
ggctcagagc	tctgaaatta	aaaagcactg	aaatagttag	atgttttcaa	acttttagaa	1860
ctcacatttt	aatcagggat	tgcatttatt	ggctaactgc	atgacatgcc	cattctacca	1920
tttaaaaaaa	aatcttaaa	cctgttaatt	ctgagaaaag	agtacagcat	ttactcttat	1980
catctagaaa	tgtaatatgc	ttcccccccg	ctttttgatg	aggaagaaga	caattggata	2040
agatgggaca	gcacttataa	tgaaataaaa	aaaaactttg	agccctctct	attccacttt	2100
agcaatcttt	ttggtaagaa	ctcttaaaag	caaaagctct	ctgaaaagat	ttgctgatta	2160
ttagtttaaa	aatcttgtaa	cactcagcag	tgctattttg	agtcacccca	gtttcctgaa	2220
agtaatgccc	agtcttctcg	aatctctctt	aatagcagaa	ccttgggtgat	tttgttggct	2280
catatgaatg	cttgtcatgg	atatgttaac	aatttagtgt	ttgacattgc	ttcctctgcc	2340
acaaagacaa	tactctgggt	acacatgtct	agaccagca	caggctgtag	gcccaggagt	2400
gactcaaagg	agtttttccc	tctttcttac	ggttcaaagg	tgaccctggg	ggtggccaga	2460
gcagtaatgc	ttgttttgat	ctcttctatg	ctcatctgct	tctcagaacc	caccggttga	2520
gtttgtgggt	aaccagcagg	caggccaaag	actgggtgct	ttcatttcat	ccttttagagg	2580
gatgaacacg	ttatttccgt	ctgatgagca	ttcggtagaa	tttttgaagt	gagattttat	2640
gaagtcaaa	gggactttac	acagatctcg	acctgctttg	aaacctagag	gtggcccttt	2700
gatttgtgcg	gttctttgcc	ctctggacaa	cttaatatatt	caagtaatcg	aataccaaact	2760
tccctgccag	cccacctgcc	ttccgccccg	cttgtgtaac	agtcctgttt	tgttgagttg	2820
ctgctattgc	actgccagtg	cagccacac	caaatcacaa	cccaagatac	tcagatagga	2880
agactcttc	cttcccagtg	actttaccaa	aggaaccccc	gccaggaccc	acatggggcc	2940
acgtgttggc	agtgggaatc	gcctgtgcag	gctggggatc	tcaggctgat	cagtaggggc	3000
cagcttttga	gccagccaag	ctgaatccca	cactccaggt	ctgtgctcaa	gagaccagat	3060

ggtgtatttc	caaatgggcc	tctctggtat	gggcaatagg	caagctcctg	gggtctgggt	3120
atgtggaaga	ttcttagtgg	atgttccgcc	tggttagctg	gttctcttca	gagaatataa	3180
agtgaatgcc	tttaggggta	gctctgaaag	agaaacccaa	caacttcatt	cctagccatg	3240
aaagtagcac	gatcatattg	tactgtattg	ttattgtaaa	atgattattt	gccatgtcat	3300
gagtaggtag	atgttttgcc	acaaatatga	aagtgtttgt	tggtcctgac	tttaagccat	3360
gaagattgag	accaataaat	agcactcaga	ggaatgaaaa	aaaaaaaaaa	agg	3413

THIS PAGE BLANK (USPTO)